

August 23, 2007

10/803.667

=> fil hcap
FILE 'HCAPLUS' ENTERED AT 16:47:30 ON 23 AUG 2007
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)


Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 23 Aug 2007 VOL 147 ISS 9
FILE LAST UPDATED: 22 Aug 2007 (20070822/ED)

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> d que 180
L1 3608 SEA FILE=REGISTRY ABB=ON PLU=ON NC4-C6/ES AND C6/ES AND N=2
AND NR=3 AND C-23
L15 12 SEA FILE=REGISTRY ABB=ON PLU=ON C2H27N2.CLO4/MF
L16 1 SEA FILE=REGISTRY ABB=ON PLU=ON L15 AND L1
L18 13 SEA FILE=REGISTRY ABB=ON PLU=ON C2H23N2S.CLO4/MF
L19 1 SEA FILE=REGISTRY ABB=ON PLU=ON L18 AND NRS=2 AND NR=3 AND
NCSC2-C6/ES AND C6/ES
L20 1 SEA FILE=REGISTRY ABB=ON PLU=ON C3H43N4S2.3BR/MF
L21 2 SEA FILE=REGISTRY ABB=ON PLU=ON C53H62N6S2.4I/MF
L25 1 SEA FILE=REGISTRY ABB=ON PLU=ON L21 NOT 177597-81-8
L26 1 SEA FILE=REGISTRY ABB=ON PLU=ON C26H31N3S.2I/MF
L32 2 SEA FILE=REGISTRY ABB=ON PLU=ON C27H32N4O6S2.2C6H15N/MF
L33 1 SEA FILE=REGISTRY ABB=ON PLU=ON C35H27BF6N3O7S2.NA/MF
L34 1 SEA FILE=REGISTRY ABB=ON PLU=ON C35H28BF6N4O7S.NA/MF
L38 STR



C-Ak
36 37
CH-CH
38 39
O-Ak
40 41
CH-Ak
42 43
O-C-Ak
44 45 46

VAR G1=H/35
VAR G2=S/O/36

August 23, 2007


10/803.667

REP G3=(1-2) 38-8 39-16
VAR G4=H/35/40
VAR G5=CH2/42
VAR G6=H/45/35

NODE ATTRIBUTES:
DEFAULT MLEVEL IS ATOM
GGCAT IS SAT AT 35
GGCAT IS SAT AT 37
GGCAT IS SAT AT 41
GGCAT IS SAT AT 46
DEFAULT ECLEVEL IS LIMITED
ECOUNT IS X3 C AT 35
ECOUNT IS X3 C AT 37
ECOUNT IS X3 C AT 41

GRAPH ATTRIBUTES:
RING(S) ARE ISOLATED OR EMBEDDED
NUMBER OF NODES IS 46

STEREO ATTRIBUTES: NONE
L40 20 SEA FILE=REGISTRY SSS FUL L38
L41 10 SEA FILE=REGISTRY ABB=ON PLU=ON L40 AND NC=2
L42 STR



O-Ak
29 30 31
Ak 32 C 33 C-G8
34 35
Ak 36 O-Ak
37 38

VAR G1=S/O/24
VAR G2=H/26
REP G3=(0-2) 27-8 28-11
VAR G4=H/30/32
VAR G7=33/34
VAR G8=36/37

NODE ATTRIBUTES:
CONNECT IS E2 RC AT 11
CONNECT IS E2 RC AT 13
CONNECT IS E2 RC AT 14
CONNECT IS E2 RC AT 24
CONNECT IS E1 RC AT 25
CONNECT IS E1 RC AT 26
CONNECT IS E2 RC AT 27
CONNECT IS E2 RC AT 28
CONNECT IS E1 RC AT 31
CONNECT IS E1 RC AT 32
CONNECT IS E2 RC AT 33
CONNECT IS E3 RC AT 34
CONNECT IS E1 RC AT 36
CONNECT IS E1 RC AT 38

DEFAULT MLEVEL IS ATOM
GGCAT IS LOC AT 26
GGCAT IS LOC AT 36
GGCAT IS LOC AT 38
DEFAULT ELEVEL IS LIMITED

GRAPH ATTRIBUTES:
RING(S) ARE ISOLATED OR EMBEDDED
NUMBER OF NODES IS 38

STEREO ATTRIBUTES: NONE
L44 95 SEA FILE=REGISTRY SSS FULL L42
L45 48 SEA FILE=REGISTRY ABB=ON PLU=ON L44 AND NC=2
L46 67 SEA FILE=REGISTRY ABB=ON PLU=ON L16 OR L19 OR L20 OR L25 OR
L26 OR L32 OR L33 OR L34 OR L41 OR L45
L47 526 SEA FILE=CAPLUS ABB=ON PLU=ON L46
L48 46411 SEA FILE=CAPLUS ABB=ON PLU=ON URINE ANALYSIS+PPT/NT/CT
L49 14072 SEA FILE=CAPLUS ABB=ON PLU=ON STAINING, BIOLOGICAL+PPT/CT
L50 1842 SEA FILE=CAPLUS ABB=ON PLU=ON STAINS, BIOLOGICAL+PPT/CT
L51 58 SEA FILE=CAPLUS ABB=ON PLU=ON L47 AND (L54 OR L55)
L52 192 SEA FILE=CAPLUS ABB=ON PLU=ON L46(L) ANST+NT/RL
L53 122 SEA FILE=CAPLUS ABB=ON PLU=ON L46(L) BIOL+NT/RL
L54 50 SEA FILE=CAPLUS ABB=ON PLU=ON L57 AND L58
L55 11 SEA FILE=CAPLUS ABB=ON PLU=ON L56 AND L59
L56 11 SEA FILE=CAPLUS ABB=ON PLU=ON L47 AND L52
L57 11 SEA FILE=CAPLUS ABB=ON PLU=ON L47 AND URIN?
L58 11 SEA FILE=CAPLUS ABB=ON PLU=ON L61 OR L62
L59 6 SEA FILE=CAPLUS ABB=ON PLU=ON L64 AND L57
L60 5 SEA FILE=CAPLUS ABB=ON PLU=ON L64 AND L58
L61 22 SEA FILE=CAPLUS ABB=ON PLU=ON L65 OR L66 OR L60
L62 29 SEA FILE=CAPLUS ABB=ON PLU=ON (L57 OR L58) AND ?BACTER?
L63 46 SEA FILE=CAPLUS ABB=ON PLU=ON L67 OR L68
L64 29 SEA FILE=CAPLUS ABB=ON PLU=ON L69 AND ?STAIN?
L65 46 SEA FILE=CAPLUS ABB=ON PLU=ON L69 OR L71
L66 4829 SEA FILE=CAPLUS ABB=ON PLU=ON SAKAI Y7/AU
L67 2302 SEA FILE=CAPLUS ABB=ON PLU=ON KAWASHIMA Y7/AU
L68 989 SEA FILE=CAPLUS ABB=ON PLU=ON INOUE J7/AU
L69 293 SEA FILE=CAPLUS ABB=ON PLU=ON IKEUCHI Y7/AU
L70 7 SEA FILE=CAPLUS ABB=ON PLU=ON (L74 OR L75 OR L76 OR L77)
L71 AND ?BACTER? AND ?STAIN?
L72 2 SEA FILE=CAPLUS ABB=ON PLU=ON L73 AND L78
L73 46 SEA FILE=CAPLUS ABB=ON PLU=ON L73 OR L79

=> d l80 ibib abs hitind hitstr tot

L80 ANSWER 1 OF 46 HCAPLUS. COPYRIGHT 2007 ACS ON STN
ACCESSION NUMBER: 2007:65371 HCAPLUS Full-text
DOCUMENT NUMBER: 146:245116
TITLE: Towards a portable microchip system with integrated thermal control and polymer waveguides for real-time PCR

AUTHOR(S): Wang, Zhenyu; Sekulovic, Andrea; Kutter, Jorg P.;
CORPORATE SOURCE: Bang, Dang D.; Wolff, Anders
MIC - Department of Micro and Nanotechnology,
Technical University of Denmark, Lyngby, Den.
SOURCE: Electrophoresis (2006), 27(24), 5051-5058
CODEN: ELCUDN; ISSN: 0173-0835
PUBLISHER: Wiley-VCH Verlag GmbH & Co. KGaA
DOCUMENT TYPE: Journal

LANGUAGE: English

AB A novel real-time PCR microchip platform with integrated thermal system and polymer waveguides has been developed. The integrated polymer optical system for real-time monitoring of PCR was fabricated in the same SU-8 layer as the PCR chamber, without addnl. masking steps. Two suitable DNA binding dyes, SYTOX Orange and TO-PRO-3, were selected and tested for the real-time PCR processes. As a model, cadp gene of Campylobacter jejuni has been amplified on the microchip. Using the integrated optical system of the real-time PCR microchip, the measured cycle threshold values of the real-time PCR performed with a dilution series of C. jejuni DNA template (2 to 200 pg/ μ L) could be quant. detected and compared with a conventional post-PCR anal. (DNA gel electrophoresis). The presented approach provided reliable real-time quant. information of the PCR amplification of the targeted gene. With the integrated optical system, the reaction dynamics at any location inside the micro reaction chamber can easily be monitored.

CC 3-1 (Biochemical Genetics)
ST Section cross-reference(s): 9, 10
IT microchip real time PCR polymer waveguide Campylobacter detection

IT Biochips
Lab-on-a-chip
Campylobacter jejuni

Optical waveguides
Temperature effects, biological
(towards a portable microchip system with integrated thermal control and polymer waveguides for real-time PCR)

IT 157199-63-8, To-PRO-3 324767-53-5, SYTOX Orange

RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)

(DNA-binding dye; towards a portable microchip system with integrated thermal control and polymer waveguides for real-time PCR)

IT 157199-63-8, To-PRO-3


RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)

(DNA-binding dye; towards a portable microchip system with integrated thermal control and polymer waveguides for real-time PCR)

RN 157199-63-8 HCAPLUS

CN Quinolinium, 4-[3-(3-methyl-2(3H)-benzothiazolylidene)-1-propen-1-yl]-1-[3-(trimethylammonio)propyl]-, iodide (1:2) (CA INDEX NAME)

(CH₂)₃-N⁺Me₃



● 2 I⁻

REFERENCE COUNT: 51 THERE ARE 51 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L80 ANSWER 2 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2007:53333 HCAPLUS Full-text
 DOCUMENT NUMBER: 146:333375
 TITLE: Binding of Intercalating and Groove-Binding Cyanine
 Dyes to Bacteriophage T5
 AUTHOR(S): Eriksson, Maja; Haerdelin, Maria; Larsson, Anette; Bergenholtz, Johan; Akerman, Björn
 CORPORATE SOURCE: Department of Chemical and Biological Engineering, Chalmers University of Technology, Goeteborg, S412 96, Swed.
 SOURCE: Journal of Physical Chemistry B (2007), 111(5), 1139-1148
 CODEN: JPCBPK; ISSN: 1520-6106
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB The interaction between four related cyanine dyes and bacteriophage T5 is investigated with fluorescence and absorption spectroscopy. The dyes, which differ in size, charge, and mode of DNA-binding, penetrate the capsid and bind the DNA inside. The rate of association decreases progressively with increasing dye size, from a few minutes for YO to more than 50 h for YOYO (at 37°). The relative affinity for the phage DNA is a factor of about 0.2 lower than for the same T5-DNA when free in solution. Comparison of groove-bound BOXTO-PRO and intercalating YO-PRO shows that the reduced affinity is not due to DNA extension but perhaps influenced by competition with other cationic DNA-binding agents inside the capsid. Although, the extent of dye binding to the phages decreases with increasing external ionic strength, the affinity relative to free DNA increases, which indicates a comparatively weak screening of electrostatic interactions inside the phage. The rate of binding increases with increasing ionic strength, reflecting an increase in effective pore size of the capsid as electrostatic interactions are screened and/or a faster diffusion of the dye through the DNA matrix inside the capsid as the DNA affinity is reduced. A combination of electron microscopy, light scattering, and linear dichroism show that the phages are intact after YO-PRO binding, whereas a small degree of capsid rupture cannot be excluded with BOXTO-PRO.

CC 9-5 (Biochemical Methods)
 Section cross-reference(s): 10
 ST Intercalating groove binding cyanine dye bacteriophage T5 assocn
 IT Molecular recognition
 (DNA; binding of intercalating and groove-binding cyanine dyes to bacteriophage T5)

IT Affinity
 Cyanine dyes
 Diffusion
 Electric screening
 Electrostatic force
 Enterobacteria phage T5
 Fluorescence
 Fluorescent indicators
 Fluorometry
 Intercalating agents
 Intercalation
 Ionic strength
 Pore size

UV and visible spectroscopy
 (binding of intercalating and groove-binding cyanine dyes to bacteriophage T5)

IT DNA
 RL: BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); BIOL (Biological study); PROC (Process)
 (binding of intercalating and groove-binding cyanine dyes to bacteriophage T5)

IT Permeability
 (capsid wall; binding of intercalating and groove-binding cyanine dyes to bacteriophage T5)

IT Virion structure
 (capsid; binding of intercalating and groove-binding cyanine dyes to bacteriophage T5)

IT Molecular association
 (dye-DNA; binding of intercalating and groove-binding cyanine dyes to bacteriophage T5)


IT 143413-85-8 143413-86-9 152068-09-2 923582-33-6, BOXTO-PRO
 RL: BUU (Biological use, unclassified); PEP (Physical, engineering or chemical process); BIOL (Biological study); PROC (Process); USES (Uses)

(binding of intercalating and groove-binding cyanine dyes to bacteriophage T5)

IT 143413-86-9
 RL: BUU (Biological use, unclassified); PEP (Physical, engineering or chemical process); BIOL (Biological study); PROC (Process); USES (Uses)

(binding of intercalating and groove-binding cyanine dyes to bacteriophage T5)

RN 143413-86-9 HCAPLUS
 CN Quinolium, 1-methyl-4-[(3-methyl-2(3H)-benzoxazolylidene)methyl]-, iodide (1:1) (CA INDEX NAME)



REFERENCE COUNT: 34 THERE ARE 34 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L80 ANSWER 3 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2007:2236 HCAPLUS Full-text
 DOCUMENT NUMBER: 146:138286
 TITLE: Reference object for detecting malfunction of particle analyzer
 INVENTOR(S): Kawate, Yasunori

PATENT ASSIGNEE(S): Symex Corporation, Japan
 SOURCE: Faming Zhuanli Shengqing Gongkai Shuomingshu, 36pp.
 CODEN: CNXHEV
 DOCUMENT TYPE: Patent
 LANGUAGE: Chinese
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO. KIND DATE APPLICATION NO. DATE
 CN 1880942 A 20061220 CN 2006-10098740 20060712
 EP 174145 A2 20070117 EP 2006-447087 20060706
 R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE,
 IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, AL,
 BA, HR, MK, YU
 JP 2007047154 A 20070222 JP 2006-189883 20060710
 PRIORITY APPL. INFO.: JP 2005-203279 A 20050712
 AB The title particle analyzer treats the target particles in the biosample by
 fluorescent staining with a certain dye, and then analyzes the stained target
 particles. The title reference object comprises a first standard particle
 treated by fluorescent staining, and a second standard particle containing
 fluorescent dye that can exhibit a certain fluorescence intensity. This
 invention also provides the method and device that uses the reference object
 to detect the abnormal parts of the particle analyzer.

CC 9-16 (Biochemical Methods)
 ST fluorescence dye emulsion ref particle fluorometry microorganism blood
 urine

IT Staining, biological
 (fluorescent; reference object for detecting malfunction of particle
 analyzer)

IT Blood analysis
 Cylinders
 Epithelium
 Erythrocyte
 Eubacteria
 Fluorescent dyes
 Fluorometry
 Leukocyte
 Light sources
 Microorganism
 Particles
 Sensors

Staining, biological
 Urine analysis


IT (reference object for detecting malfunction of particle analyzer)
 514-73-8, NK-136 36536-22-8, NK-529 189148-50-3
 RL: ARG (Analytical reagent use); ANST (Analytical
 study); USES (Uses)

IT (reference object for detecting malfunction of particle analyzer)
 189148-50-3
 RL: ARG (Analytical reagent use); ANST (Analytical
 study); USES (Uses)

RN (reference object for detecting malfunction of particle analyzer)
 CN Quinolinium, 1-(2-hydroxyethyl)-4-[3-(3-methyl-2(3H)-benzothiazolylidene)-
 1-propenyl]-, tetrafluoroborate (1-) (9CI) (CA INDEX NAME)

CM 1

CRN 189148-49-0
 CMF C22 H21 N2 O S



CM 2

CRN 14874-70-5
 CMF B P4
 CCI CCS



L80 ANSWER 4 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2006:1207557 HCAPLUS Full-text
 DOCUMENT NUMBER: 145:495830
 TITLE: Disinfection of biological fluids using asymmetric
 cyanine dyes

INVENTOR(S): Wagner, Stephen J.
 PATENT ASSIGNEE(S): USA
 SOURCE: U.S. Pat. Appl. Publ., 14pp.
 CODEN: USXXCO

DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO. KIND DATE APPLICATION NO. DATE
 US 2006257844 A1 20061116 US 2006-430848 20060510
 PRIORITY APPL. INFO.: US 2005-680502P P. 20050513
 OTHER SOURCE(S): MARPAT 145:495830

AB Asym. cyanine dyes that bind nucleic acid but not red blood cell membrane, and
 function as photosensitizers when rigidly bound but not when free in solution
 are provided. Unbound dye thus causes minimal oxidative damage. The dyes do

not substantially accumulate in red blood cells, thereby minimizing hemolysis due to oxidative damage. Biol. fluids, such as blood and blood products can be disinfected by mixing the fluid with these asym. cyanine dyes that binds to nucleic acid, irradiating the mixture, recovering clin. significant components from the biol. fluid and/or assaying the fluid for pathogens. Thus, various viruses and bacteria were photoactivated in a red blood cell preparation using Thiazole Orange. For example, Thiazole Orange phototreatment using 80 µm dye and 7.4 J/cm² of cool white light inactivated 5.1 to 7.4 log₁₀ of tested envelope viruses. In addition, > 6.3 log₁₀ of intracellular HIV was photoactivated.

INCL 435002000; 435032000; 435031000

CC 63-8 (Pharmaceuticals)

IT Animal virus

Blood

Blood cell

Blood products

Cyanine dyes

Disinfectants

Erythrocyte

Eubacteria

Human

Human immunodeficiency virus 1

Light

Photosensitizers, pharmaceutical

Sterilization and Disinfection

(disinfection of biol. fluids using asym. cyanine dyes followed by irradiation)

IT 107091-89-4, Thiazole Orange

RL: THU (Therapeutic use); BIOL (Biological study);

USES (Uses)

(disinfection of biol. fluids using asym. cyanine dyes followed by irradiation)

(disinfection of biol. fluids using asym. cyanine dyes followed by irradiation)

IT 107091-89-4, Thiazole Orange

RL: THU (Therapeutic use); BIOL (Biological study);

USES (Uses)

(disinfection of biol. fluids using asym. cyanine dyes followed by irradiation)

(disinfection of biol. fluids using asym. cyanine dyes followed by irradiation)

IT 107091-89-4, Thiazole Orange

RL: THU (Therapeutic use); BIOL (Biological study);

USES (Uses)

(disinfection of biol. fluids using asym. cyanine dyes followed by irradiation)

(disinfection of biol. fluids using asym. cyanine dyes followed by irradiation)

IT 107091-89-4, Thiazole Orange

RL: THU (Therapeutic use); BIOL (Biological study);

USES (Uses)

(disinfection of biol. fluids using asym. cyanine dyes followed by irradiation)

(disinfection of biol. fluids using asym. cyanine dyes followed by irradiation)

IT 107091-89-4, Thiazole Orange

RL: THU (Therapeutic use); BIOL (Biological study);

USES (Uses)

(disinfection of biol. fluids using asym. cyanine dyes followed by irradiation)

(disinfection of biol. fluids using asym. cyanine dyes followed by irradiation)

IT 107091-89-4, Thiazole Orange

RL: THU (Therapeutic use); BIOL (Biological study);

USES (Uses)

(disinfection of biol. fluids using asym. cyanine dyes followed by irradiation)

(disinfection of biol. fluids using asym. cyanine dyes followed by irradiation)

IT 107091-89-4, Thiazole Orange

RL: THU (Therapeutic use); BIOL (Biological study);

USES (Uses)

(disinfection of biol. fluids using asym. cyanine dyes followed by irradiation)

(disinfection of biol. fluids using asym. cyanine dyes followed by irradiation)

IT 107091-89-4, Thiazole Orange

RL: THU (Therapeutic use); BIOL (Biological study);

USES (Uses)

(disinfection of biol. fluids using asym. cyanine dyes followed by irradiation)

(disinfection of biol. fluids using asym. cyanine dyes followed by irradiation)

IT 107091-89-4, Thiazole Orange

RL: THU (Therapeutic use); BIOL (Biological study);

USES (Uses)

(disinfection of biol. fluids using asym. cyanine dyes followed by irradiation)

(disinfection of biol. fluids using asym. cyanine dyes followed by irradiation)

IT 107091-89-4, Thiazole Orange

RL: THU (Therapeutic use); BIOL (Biological study);

USES (Uses)

(disinfection of biol. fluids using asym. cyanine dyes followed by irradiation)

(disinfection of biol. fluids using asym. cyanine dyes followed by irradiation)

IT 107091-89-4, Thiazole Orange

RL: THU (Therapeutic use); BIOL (Biological study);

USES (Uses)

(disinfection of biol. fluids using asym. cyanine dyes followed by irradiation)

(disinfection of biol. fluids using asym. cyanine dyes followed by irradiation)

IT 107091-89-4, Thiazole Orange

RL: THU (Therapeutic use); BIOL (Biological study);

USES (Uses)

(disinfection of biol. fluids using asym. cyanine dyes followed by irradiation)

(disinfection of biol. fluids using asym. cyanine dyes followed by irradiation)


IT 107091-89-4, Thiazole Orange

RL: THU (Therapeutic use); BIOL (Biological study);

USES (Uses)

(disinfection of biol. fluids using asym. cyanine dyes followed by irradiation)

(disinfection of biol. fluids using asym. cyanine dyes followed by irradiation)



CM 2

CRN 16722-51-3

CMF C7 H7 O3 S



L80 ANSWER 5 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 2006:1123770 HCAPLUS Full-text
DOCUMENT NUMBER: 145:451262
TITLE: Multiplex microparticle system for microarrays including fluorescence dye-bound polymer beads
INVENTOR(S): Mehrpouyan, Majid; Recktenwald, Diether J.; Varro, Rudolf
PATENT ASSIGNEE(S): Becton, Dickinson and Company, USA
SOURCE: U.S. Pat. Appl. Publ., 12pp.
CODEN: USXXCO
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|---|------|----------|-----------------|----------|
| US 2006240411 | A1 | 20061026 | US 2006-404348 | 20060414 |
| WO 2006115870 | A2 | 20061102 | WO 2006-US14361 | 20060414 |
| WO 2006115870 | A3 | 20070719 | | |
| W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW | | | | |
| RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GN, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA | | | | |

PRIORITY APPLN. INFO.:
AB Arrays of microparticle populations, each population labeled with a single fluorescent dye, are provided for use in multiplex assays. The populations form a virtual multidimensional array wherein each microparticle is identified by fluorescence intensity in two different fluorescence detection channels. The arrays are useful in a variety of assays, including multiplex, multi-analyte assays for the simultaneous detection of two or more analytes by, for example, flow cytometry, and a labeling reagents in, for example, microscopy. The use of singly-dyed microparticles to form multidimensional arrays greatly simplifies the creation of multiplex assays.

INCL 435005000; 435006000; 435007310; 435007320; 435018000
CC 9-1 (Biochemical Methods)

10/803,667

August 23, 2007

August 23, 2007

10/803,667

Section cross-reference(s): 3, 4

IT Algae

Animal tissue

Cell

Environmental analysis

Eubacteria

Fluorescence microscopy

Fluorescent dyes

Fluorometry

Fungi

Microarray technology

Microorganism

Microparticles

Parasite

Pathogen

Virus

(multiplex microparticle system for microarrays including fluorescence dye-bound polymer beads)

IT 24796-94-9, Oxazine 725 76433-27-7, LDS 730 76433-29-9

, LDS 751 89872-07-1, LDS 750 154530-43-5, LDS 765 913169-98-9, ABS 643

RL: ARG (Analytical reagent use); PRP (Properties); ANST

(Analytical study; USES (Uses))

(multiplex microparticle system for microarrays including fluorescence dye-bound polymer beads)

IT 76433-27-7, LDS 730 76433-29-9, LDS 751

RL: ARG (Analytical reagent use); PRP (Properties); ANST

(Analytical study; USES (Uses))

(multiplex microparticle system for microarrays including fluorescence dye-bound polymer beads)


RN 76433-27-7 HCAPLUS

CN 3H-Indolium, 2-[4-[4-(dimethylamino)phenyl]-1,3-butadienyl]-1,3,3-trimethyl-, perchlorate (9CI) (CA INDEX NAME)

CM 1

CRN 76433-26-6

CMF C23 H27 N2



CM 2

CRN 14797-73-0

CMF C1 O4



RN 76433-29-9 HCAPLUS

CN Benzothiazolium, 2-[4-[4-(dimethylamino)phenyl]-1,3-butadienyl]-3-ethyl-, perchlorate (1:1) (CA INDEX NAME)

CM 1

CRN 76433-28-8

CMF C21 H23 N2 S



CM 2

CRN 14797-73-0

CMF C1 O4



L80 ANSWER 6 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 2006:630000 HCAPLUS Full-text
DOCUMENT NUMBER: 145:79328
TITLE: A method for diagnosing and monitoring cellular reservoirs of disease

INVENTOR(S): Scott, Lesley Erica

PATENT ASSIGNER(S): University of the Witwatersrand, Johannesburg, S. Afr.

SOURCE: PCT Int. Appl., 39 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO. KIND DATE APPLICATION NO. DATE

WO 2006067572 A2 20060629 WO 2005-1B3738 20051212
 WO 2006067572 A3 20060810

W: AB, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DS, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RU, SC, SD, SE, SG, SK, SL, SM, ST, TJ, TW, TN, TR, TT, TZ, UA, UG, US, VZ, VE, VN, YU, ZA, ZM, ZW

RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM

PRIORITY APPLN. INFO.:
 AB The invention provides an assay for diagnosing and/or monitoring a viral infection or disease in a patient, the assay including the steps of mixing a sample of leukocytes with a fluorescent cell membrane-permeable dye which stains RNA or both DNA and RNA within the leukocytes; identifying from all the leukocytes at least two of the three major sub-populations of leukocytes selected from the group consisting of monocytes, granulocytes and lymphocytes; determining the fluorescence intensity for each of the identified sub-populations; and comparing the fluorescence intensity of at least two cell sub-populations to each other to obtain at least one of the following ratios: monocytes to granulocytes, monocytes to lymphocytes, and granulocytes to lymphocytes. The viral infection may be HIV and the disease may be AIDS. The invention also provides a method of monitoring the cellular viral, parasitic or bacterial reservoir of a patient with a viral or bacterial infection by the steps described above. A kit for performing the assay or method is also provided. Red blood cells of blood samples from HIV neg. and HIV pos. patients were lysed using Immunoprep reagent and lysed samples were treated with thiazole orange. The samples were then analyzed on an XL MCL flow cytometer and the mean fluorescent intensity in the FL1 channel for each region was measured. The monocyte, granulocyte and lymphocyte ratios were determined. The ratios were different for HIV pos. patients compared with HIV neg. patients. The monocytes from HIV pos. patients had increased mean fluorescent intensity.

9-5 (Biochemical Methods)
 Section cross-reference(s): 10, 14, 15
 diagnosis cellular reservoir disease; leukocyte RNA staining subpopulation fluorescence viral infection assay; kit assay disease reservoir leukocyte RNA fluorescence; HIV monocyte mean fluorescent intensity thiazole orange staining

IT Anti-AIDS agents
 Antiviral agents
 (HIV reservoir monitoring in relation to treatment with; assay for diagnosing and monitoring cellular reservoirs of disease and assay kit containing fluorescent RNA-staining cell membrane-permeable dye)

IT CD14 (antigen)
 CD38 (antigen)
 FcγRIII receptors
 RL: ANT (Analyte); BSU (Biological study, unclassified); DGN (Diagnostic use); ANST (Analytical study); BIOL (Biological study); USES (Uses) (as cell activation marker for phenotyping; assay for diagnosing and monitoring cellular reservoirs of disease and assay kit containing fluorescent RNA-staining cell membrane-permeable dye)

IT Tuberculosis
 (as other disease, addnl. monitoring of; assay for diagnosing and monitoring cellular reservoirs of disease and assay kit containing fluorescent RNA-staining cell membrane-permeable dye)

IT AIDS (disease)
 Animal tissue culture
 Blood analysis
 Cell membrane
 Computer program
 Development, mammalian postnatal
 Diagnosis
 Disease, animal
 Electric impedance
 Flow cytometry
 Fluorescence
 Fluorescent dyes
 Fluorometry
 Human
 Human immunodeficiency virus
 Human immunodeficiency virus 1
 Lymphocyte
 Monocyte
 Polymorphonuclear leukocyte
 Prognosis
 Test kits
 (assay for diagnosing and monitoring cellular reservoirs of disease and assay kit containing fluorescent RNA-staining cell membrane-permeable dye)

IT DNA
 RNA
 RL: ANT (Analyte); BSU (Biological study, unclassified); DGN (Diagnostic use); ANST (Analytical study); BIOL (Biological study); USES (Uses) (assay for diagnosing and monitoring cellular reservoirs of disease and assay kit containing fluorescent RNA-staining cell membrane-permeable dye)

IT CD4 (antigen)
 RL: ANT (Analyte); BSU (Biological study, unclassified); DGN (Diagnostic use); ANST (Analytical study); BIOL (Biological study); USES (Uses) (assay further including count of; assay for diagnosing and monitoring cellular reservoirs of disease and assay kit containing fluorescent RNA-staining cell membrane-permeable dye)

IT Fluorometers
 (assay using; assay for diagnosing and monitoring cellular reservoirs of disease and assay kit containing fluorescent RNA-staining cell membrane-permeable dye)

IT Infection
 (bacterial; assay for diagnosing and monitoring cellular reservoirs of disease and assay kit containing fluorescent RNA-staining cell membrane-permeable dye)

IT Spheres
 (beads, kit containing reagents of; assay for diagnosing and monitoring cellular reservoirs of disease and assay kit containing fluorescent RNA-staining cell membrane-permeable dye)

IT Infection
 (co-infection, addnl. monitoring of; assay for diagnosing and monitoring cellular reservoirs of disease and assay kit containing fluorescent RNA-staining cell membrane-permeable dye)

IT Therapy
 (determination of patient response to; assay for diagnosing and monitoring cellular reservoirs of disease and assay kit containing fluorescent RNA-staining cell membrane-permeable dye)

IT Antibodies and immunoglobulins
 RL: ARG (Analytical reagent use); BSU (Biological study, unclassified); DGN (Diagnostic use); ANST (Analytical study); BIOL (Biological study);

USES (Uses)
(fluorescent, for CD4 count, assay further including, assay for
diagnosing and monitoring cellular reservoirs of disease and assay kit
containing fluorescent RNA-staining cell membrane-permeable dye)

IT Staining, biological
(fluorescent; assay for diagnosing and monitoring cellular reservoirs
of disease and assay kit containing fluorescent RNA-staining cell
membrane-permeable dye)

IT Immunoassay
(for CD14/CD16 immunophenotyping; assay for diagnosing and monitoring
cellular reservoirs of disease and assay kit containing fluorescent RNA-
staining cell membrane-permeable dye)

IT Blood
(hematol. analyzer, assay using; assay for diagnosing and monitoring
cellular reservoirs of disease and assay kit containing fluorescent RNA-
staining cell membrane-permeable dye)

IT Development, mammalian postnatal
(infant; assay for diagnosing and monitoring cellular reservoirs of
disease and assay kit containing fluorescent RNA-staining cell
membrane-permeable dye)

IT Cytolysis
(kit containing agent for red blood cell; assay for diagnosing and
monitoring cellular reservoirs of disease and assay kit containing
fluorescent RNA-staining cell membrane-permeable dye)

IT Culture media
Stabilizing agents
(kit containing; assay for diagnosing and monitoring cellular reservoirs of
disease and assay kit containing fluorescent RNA-staining cell
membrane-permeable dye)

IT Computers
(kit including instructions readable by; assay for diagnosing and
monitoring cellular reservoirs of disease and assay kit containing
fluorescent RNA-staining cell membrane-permeable dye)

IT UV and visible spectroscopy
(light-scattering; assay for diagnosing and monitoring cellular
reservoirs of disease and assay kit containing fluorescent RNA-
staining cell membrane-permeable dye)

IT Erythrocyte
(lysis in blood sample for anal. of leukocytes; assay for diagnosing
and monitoring cellular reservoirs of disease and assay kit containing
fluorescent RNA-staining cell membrane-permeable dye)

IT Cell activation
(markers, for phenotyping; assay for diagnosing and monitoring cellular
reservoirs of disease and assay kit containing fluorescent RNA-
staining cell membrane-permeable dye)

IT Samples
(of leukocytes; assay for diagnosing and monitoring cellular reservoirs
of disease and assay kit containing fluorescent RNA-staining cell
membrane-permeable dye)

IT gag proteins
RL: ANT (Analyte); BSU (Biological study, unclassified); DGN (Diagnostic
use); ANST (Analytical study); BIOL (Biological study); USES (Uses)
(p24gag, as cell activation marker for phenotyping; assay for
diagnosing and monitoring cellular reservoirs of disease and assay kit
containing fluorescent RNA-staining cell membrane-permeable dye)

IT Infection
(parasitic; assay for diagnosing and monitoring cellular reservoirs of
disease and assay kit containing fluorescent RNA-staining cell
membrane-permeable dye)

IT Biological transport

(permeation, of fluorescent dye through cell membrane; assay for
diagnosing and monitoring cellular reservoirs of disease and assay kit
containing fluorescent RNA-staining cell membrane-permeable dye)

IT Leukocyte
(sample of; assay for diagnosing and monitoring cellular reservoirs of
disease and assay kit containing fluorescent RNA-staining cell
membrane-permeable dye)

IT Phenotypes
(set of cell membrane markers or intracellular markers for determining;
assay for diagnosing and monitoring cellular reservoirs of disease and assay
kit containing fluorescent RNA-staining cell membrane-permeable
dye)

IT Phycocerythrins
RL: ARG (Analytical reagent use); BSU (Biological study, unclassified);
DGN (Diagnostic use); ANST (Analytical study); BIOL (Biological study);
USES (Uses)
(thiazole orange nucleic acid binding dye in combination with
CD4-binding; assay for diagnosing and monitoring cellular reservoirs of
disease and assay kit containing fluorescent RNA-staining cell
membrane-permeable dye)


IT Infection
(viral; assay for diagnosing and monitoring cellular reservoirs of
disease and assay kit containing fluorescent RNA-staining cell
membrane-permeable dye)

IT 107091-89-4, Thiazole orange
RL: ARG (Analytical reagent use); BSU (Biological study,
unclassified); DGN (Diagnostic use); ANST (Analytical
study); BIOL (Biological study); USES (Uses)
(assay for diagnosing and monitoring cellular reservoirs of disease and
assay kit containing fluorescent RNA-staining cell
membrane-permeable dye)

IT 107091-89-4, Thiazole orange
RL: ARG (Analytical reagent use); BSU (Biological study,
unclassified); DGN (Diagnostic use); ANST (Analytical
study); BIOL (Biological study); USES (Uses)
(assay for diagnosing and monitoring cellular reservoirs of disease and
assay kit containing fluorescent RNA-staining cell
membrane-permeable dye)

RN 107091-89-4 HCAPLUS
CN Quinolinium, 1-methyl-4-[(3-methyl-2(3H)-benzothiazolylidene)methyl]-,
4-methylbenzenesulfonate (1:1) (CA INDEX NAME)

CM 1
CRN 24144-08-9
CMF C19 H17 N2 S



10/803.667

August 23, 2007

August 23, 2007

10/803.667

143413-85-8, YOYO-1 154757-99-0, POPO-3 156312-20-8, YOYO-3
157199-63-8, TOPRO3 163795-75-3, SYBR Green I
166196-17-4, TOYO-3 169454-13-1, BOBO-1 169454-15-3, POPO-1
169454-17-5, BOBO-3 177571-06-1, PicoGreen 194100-76-0, SYTOX Green
305802-06-6, LOLO-1
RL: ARG (Analytical reagent use); BUU (Biological use,
unclassified); ANST (Analytical study); BIOL
(Biological study); USES (Uses)
(methods for detection of pathogens in red blood cells)
IT 107091-89-4, Thiazole orange 157199-63-8, TOPRO3
166196-17-4, TOYO-3
RL: ARG (Analytical reagent use); BUU (Biological use,
unclassified); ANST (Analytical study); BIOL
(Biological study); USES (Uses)
(methods for detection of pathogens in red blood cells)
RN 107091-89-4 HCAPLUS
CN Quinolinium, 1-methyl-4-((3-methyl-2(3H)-benzothiazolylidene)methyl)-,
4-methylbenzenesulfonate (1:1) (CA INDEX NAME)

L80 ANSWER 7 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 2006:273965 HCAPLUS Full-text
DOCUMENT NUMBER: 144:307966
TITLE: Methods for detection of pathogens in red blood cells
INVENTOR(S): Vannier, Edoard
PATENT ASSIGNEE(S): New England Medical Center Hospitals, Inc., USA
SOURCE: PCT Int. Appl., 47 pp.
CODEN: PIXXD2



DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|------------------------|--|----------|-----------------|------------|
| WO 2006031544 | A2 | 20060323 | WO 2005-US31793 | 20050909 |
| WO 2006031544 | A3 | 20070322 | | |
| W: | AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, GU, HD, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW | | | |
| RW: | AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TC, TG, BM, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM | | | |
| US 200603185 | A1 | 20060323 | US 2005-223599 | 20050909 |
| PRIORITY APPLN. INFO.: | | | US 2004-608825P | P 20040909 |
| AB | The present invention relates to methods for diagnosing a parasitic microorganism that resides in red blood cells, such as Babesia microti. | | | |
| CC | 9-16 (Biochemical Methods) | | | |
| IT | Section cross-reference(s): 10, 14 | | | |
| IT | Stains, biological (dimeric cyanine nucleic acid; methods for detection of pathogens in red blood cells) | | | |
| IT | Stains, biological (fluorescent, nucleic acid; methods for detection of pathogens in red blood cells) | | | |
| IT | 2535-16-4, Propidium iodide 107091-89-4, Thiazole orange | | | |

17


CM 2

CRN 16722-51-3
CMF C7 H7 O3 S



CM 2

CRN 16722-51-3
CMF C7 H7 O3 S



RN 157199-63-8 HCAPLUS
CN Quinolinium, 4-[3-(3-methyl-2(3H)-benzothiazolylidene)-1-propen-1-yl]-1-[3-(trimethylammonio)propyl]-, iodide (1:2) (CA INDEX NAME)


18

(CH₂)₃-N⁺Me₃

RN 166196-17-4 HCAPLUS

CN Quinolinium, 1,1'-[1,3-propanediyl]bis[(dimethyliminio)-3,1-propanediyl]bis[4-(3-methyl-2(3H)-benzothiazolylidene)-1-propen-1-yl]-, iodide (1:4) (CA INDEX NAME)

PAGE 1-A



LS0 ANSWER 8 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 2006:233900 HCAPLUS Full-text

DOCUMENT NUMBER: 144:288928

TITLE: Microorganism sterilization treatment effect-measuring method using two kinds of cell growth activity information

INVENTOR(S): Oda, Yasumasa; Sakata, Takashi

PATENT ASSIGNEE(S): Sysmex Co., Ltd., Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 12 pp.

CODEN: JKXXAF

DOCUMENT TYPE: Patent


LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|---|------|----------|-----------------|----------|
| JP 2006067974 | A | 20060316 | JP 2004-258723 | 20040906 |
| PRIORITY APPLN. INFO.: AB A method is provided for rapidly and accurately measuring the sterilization treatment effect on microorganism (e.g., bacillus). The method comprises elec. or optically measuring two kinds of growth activity information on the microorganism contained in a sample which has been treated for sterilization and cultured for a specified time, and calculating the microorganism number in a specified region (e.g., spore region, germination region, nutrition-type region) divided in a two-dimensional distribution diagram formed based on the two kinds of growth activity information. | | | JP 2004-258723 | 20040906 |
| CC 9-5 (Biochemical Methods) | | | | |
| IT Section cross-reference(s): 10 | | | | |
| IT Staining, biological (fluorescent; microorganism sterilization treatment effect-measuring method using two kinds of cell growth activity information) | | | | |
| IT Bacillus (bacteri. genus) Dimension Flow cytometry Fluorometry Germination Growth, microbial Microorganism | | | | |


Nutrition, microbial
 Sterilization and Disinfection
 (microorganism sterilization treatment effect-measuring method using two kinds of cell growth activity information)
 IT 189148-51-4
 RL: EUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)
 (microorganism sterilization treatment effect-measuring method using two kinds of cell growth activity information)
 IT 189148-51-4
 RL: EUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)
 (microorganism sterilization treatment effect-measuring method using two kinds of cell growth activity information)
 RN 189148-51-4 HCAPLUS
 CN Quinolinium, 1-(2,3-dihydroxypropyl)-4-[3-(3-methyl-2(3H)-benzothiazolylidene)-1-propenyl]-, bromide (9CI) (CA INDEX NAME)



L80 ANSWER 9 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2006:180959 HCAPLUS Full-Text
 DOCUMENT NUMBER: 144:376148
 TITLE: Thiazole orange, a DNA-binding photosensitizer with flexible structure, can inactivate pathogens in red blood cell suspensions while maintaining red cell storage properties
 AUTHOR(S): Skripchenko, Andrey; Wagner, Stephen J.; Thompson-Montgomery, Dedee; Awatefe, Helen
 CORPORATE SOURCE: Holland Laboratory, Blood Components Development, American Red Cross Biomedical Services, Rockville, MD, USA
 SOURCE: Transfusion (Walden, MA, United States) (2006), 46(2), 213-219
 CODEN: TRANAT; ISSN: 0041-1132
 PUBLISHER: Blackwell Publishing, Inc.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB Development of a robust pathogen reduction system for red cells (RBCs) utilizing photosensitive dyes was constrained by hemolysis, usually mediated by reactive oxygen species emanating from dye free in solution as well as dye


bound to the RBC membrane. The RBC binding properties of thiazole orange (TO), a flexible nucleic acid intercalating cyanine dye that predominantly acts as a photosensitizer only when bound, were assessed along with its virucidal, bactericidal, and light-induced hemolytic activities. Leukodepleted 20% hematocrit RBCs suspended in Erythrosol (RAS-2) were oxygenated, inoculated with test organisms, incubated with TO, and illuminated. Control and treated samples were analyzed by appropriate assay. Identically prepared, but uncontaminated samples were phototreated, concentrated to 45% hematocrit, and assayed for potassium leakage, hemolysis, and ATP during storage. Approx. 21 percent TO bound to RBCs. Phototreatment inactivated from 5.4 to 7.1 log10 of 5 tested viruses and from 2.3 to greater than 7.0 log10 of 8 tested bacteria. Phototreated RBCs exhibited only slightly increased hemolysis, moderately elevated potassium efflux, and similar levels of ATP compared to controls. TO can photoinactivate several model viruses and pathogens in RBCs under conditions that produce limited hemolysis without the addition of quenchers or competitive inhibitors.
 CC 63-3 (Pharmaceuticals)
 IT Section cross-reference(s): 8
 Antibacterial agents
 Antiviral agents
 Blood preservation
 Blood products
 Erythrocyte
 Light
 Pathogen
 Photodynamic action
 Photodynamic therapy
 Photosensitizers, pharmaceutical
 (thiazole orange photoinactivating pathogens in red blood cell suspensions while maintaining red cell storage properties)
 IT 107091-89-4, Thiazole orange
 RL: THU (Therapeutic use); BIOL (Biological study);
 USES (Uses)
 (thiazole orange photoinactivating pathogens in red blood cell suspensions while maintaining red cell storage properties)
 IT 107091-89-4, Thiazole orange
 RL: THU (Therapeutic use); BIOL (Biological study);
 USES (Uses)
 (thiazole orange photoinactivating pathogens in red blood cell suspensions while maintaining red cell storage properties)
 RN 107091-89-4 HCAPLUS
 CN Quinolinium, 1-methyl-4-[(3-methyl-2(3H)-benzothiazolylidene)methyl]-, 4-methylbenzenesulfonate (1:1) (CA INDEX NAME)

CM 1
 CRN 24144-08-9
 CMF C19 H17 N2 S



CM 2

CRN 16722-51-3
CMF C7 H7 O3 S



REFERENCE COUNT:

35 THERE ARE 35 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L80 ANSWER 10 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 2006:100431 HCAPLUS Full-text
DOCUMENT NUMBER: 144:187496
TITLE: Method and device for characterization of the cellular
components of a biological fluid

INVENTOR(S):

PATENT ASSIGNEE(S):

SOURCE:

Fr. Demande, 34 pp.

CODEN: FRXXEL

Patent

French

DOCUMENT TYPE:

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|---|------|----------|-----------------|----------|
| FR 2873813 | A1 | 20060203 | FR 2004-8431 | 20040730 |
| FR 2873813 | B1 | 20061117 | | |
| CA 2576753 | A1 | 20060309 | CA 2005-2576753 | 20050706 |
| WO 2006024716 | A1 | 20060309 | WO 2005-FR1740 | 20050706 |
| W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, GU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NZ, NA, NG, NI, NO, OM, PH, PL, PT, RO, RU, SC, SE, SG, SK, SL, SM, SY, TJ, TW, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW | | | | |
| RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BI, | | | | |


CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
EP 1771718 A1 20070411 EP 2005-788740 20050706
R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR
CN 1993610 A 20070704 CN 2005-80025806 20050706
FR 2004-8431 A 20040730
WO 2005-FR1740 W 20050706
PRIORITY APPLN. INFO.:
AB A method for differentiation and counting of the cellular components present in a sample of a biol. liquid includes a primary stage of cytol. anal. classically implemented by flow cytom. apparatus to obtain an ensemble of primary results allowing a differentiation and a counting of the whole of the cellular components of the sample as various populations; and a complementary stage of cytol. anal. of a particular type of cellular components, function of an identified cellular characteristic, to obtain complementary results allowing a differentiation and a counting of at least a population or cellular subpopulation of the sample for the identification of this cellular characteristic. The invention is useful in particular for hematol. anal.

9-1 (Biochemical Methods)
CC Section cross-reference(s): 15
IT Animal cell
Apparatus
Basophil
Blood
Blood analysis
Body fluid
Bone marrow
Cerebrospinal fluid
Colored materials
Colorimetry
Diagnosis
Diffusion
Electric impedance
Electric resistance
Electrodes
Eosinophil
Erythrocyte
Flow cytometry
Fluorescence
Fluorescent substances
Fluorometry
Hematopoietic precursor cell
IR radiation
Lasers
Leukocyte
Lymphocyte
Monocyte
Neutrophil
Optical absorption
Optical diffraction
Optical transmission
Optics
Platelet (blood)
Pleural fluid
Synovial fluid
UV radiation
Urine analysis
(device and method for characterization of cellular components of biol. fluid)

- IT 107091-89-4, Thiazole orange 140876-43-3, PC 5
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)
(device and method for characterization of cellular components of biol. fluid)
- IT 107091-89-4, Thiazole orange
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)
(fluid)
- RN 107091-89-4 HCAPLUS
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)
(fluid)
- CN Quinolium, 1-methyl-4-[(3-methyl-2(3H)-benzothiazolylidene)methyl]-, 4-methylbenzenesulfonate (1:1) (CA INDEX NAME)


CM 1

CRN 24144-08-9
CMF C19 H17 N2 S



CM 2

CRN 16722-51-3
CMF C7 H7 O3 S



REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L60 ANSWER 11 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 2005:1234461 HCAPLUS Full-text
DOCUMENT NUMBER: 145:3492
TITLE: Implementation of accurate and fast DNA cytometry by confocal microscopy in 3D
AUTHOR(S): Ploeger, Lennert S.; Huismans, Andre; van der Gugten, Jurryt; van der Giesen, Dionne M.; Belien, Jeroen A. M.; Abbaker, Abdelhadi Y.; Dullens, Hub F. J.; Grizzle, William; Poulin, Neal M.; Meijer, Gerrit A.;

- CORPORATE SOURCE: van Diest, Paul J.
Department of Pathology, University Medical Center
Utrecht, Utrecht, 3508 GA, Neth.
Cellular Oncology (2005), 27(4), 225-230
CODEN: COENCD; ISSN: 1570-5870
- SOURCE: IOS Press
PUBLISHER: Journal
DOCUMENT TYPE: English
LANGUAGE: English
- AB DNA cytometry is a powerful method for measuring genomic instability. Standard approaches that measure DNA content of isolated cells may induce selection bias and do not allow interpretation of genomic instability in the context of the tissue. Confocal Laser Scanning Microscopy (CLSM) provides the opportunity to perform 3D DNA content measurements on intact cells in thick histol. sections. Because the technique is tech. challenging and time consuming, only a small number of usually manually selected nuclei were analyzed in different studies, not allowing wide clin. evaluation. The aim of this study was to describe the conditions for accurate and fast 3D CLSM cytometry with a min. of user interaction to arrive at sufficient throughput for pilot clin. applications. Nuclear DNA was stained in 14 µm thick tissue sections of normal liver and adrenal stained with either YOYO-1 iodide or TO-PRO-3 iodide. Different pre-treatment strategies were evaluated: boiling in citrate buffer (pH 6.0) followed by RNase application for 1 or 18 h, or hydrolysis. The image stacks obtained with CLSM at microscope magnifications of +40 or +100 were analyzed off-line using inhouse developed software for semi-automated 3D fluorescence quantitation. To avoid sectioned nuclei, the top and bottom of the stacks were identified from ZX and YZ projections. As a measure of histogram quality, the coefficient of variation (CV) of the diploid peak was assessed. The lowest CV (10.3%) was achieved with a protocol without boiling, with 1 h RNase treatment and TO-PRO-3 iodide staining, and a final image recording at +60 or +100 magnifications. A sample size of 300 nuclei was generally achievable. By filtering the set of automatically segmented nuclei based on volume, size and shape, followed by interactive removal of the few remaining faulty objects, a single measurement was completely analyzed in approx. 3 h. The described methodol. allows to obtain a largely unbiased sample of nuclei in thick tissue sections using 3D DNA cytometry by confocal laser scanning microscopy within an acceptable time frame for pilot clin. applications, and with a CV small enough to resolve smaller near diploid stemlines. This provides a suitable method for 3D DNA ploidy assessment of selected rare cells based on morphol. characteristics and of clin. samples that are too small to prepare adequate cell suspensions.

CC

IT

9-4 (Biochemical Methods)
Animal tissue
Boiling
Cell nucleus
Confocal laser scanning microscopy
Cytometry
Dimension
Human
Imaging
Solvolytic
Staining, biological
Implementation of accurate and fast DNA cytometry by confocal microscopy in 3D

IT


143413-85-8, YOYO-1 157199-63-8, TO-PRO-3 iodide
RL: APG (Analytical reagent use); BSU (Biological study, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)
(implementation of accurate and fast DNA cytometry by confocal microscopy in 3D)

IT

157199-63-8, TO-PRO-3 iodide

RL: ARG (Analytical reagent use); BSU (Biological study, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses) (Implementation of accurate and fast DNA cytometry by confocal microscopy in 3D)

RN 157199-63-8 HCAPLUS
CN Quinolinium, 4-[3-(3-methyl-2(3H)-benzothiazolylidene)-1-propen-1-yl]-1-[3-(trimethylammonio)propyl]-, iodide (1:2) (CA INDEX NAME)



● 2 I⁻

REFERENCE COUNT:

15 THERE ARE 15 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L80 ANSWER 12 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 2005:962309 HCAPLUS Full-text
DOCUMENT NUMBER: 143:246739

TITLE: FATT-CTL (fluorescent-antigen-transfected target -- cytotoxic T lymphocyte) assay, nucleic acids and kits to detect antigen-specific cytolytic activity for immunity assessing and drug screening

INVENTOR(S):

Gruters, Robertus Antonius; Van Baalen, Carel
Adrianus; Rimmelzwaan, Guustaaf Frank; Osterhaus, Albertus Dominicus Marcellinus Erasmus
Erasmus Universiteit Rotterdam, Neth.

PATENT ASSIGNEE(S):

PCT Int. Appl., 67 pp.

CODEN: PFXD2

PATENT

English

1

FAMILY ACC. NUM. COUNT:

PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|---------------|--|----------|-----------------|----------|
| WO 2005080991 | A1 | 20050901 | WO 2005-NL119 | 20050218 |
| W: | AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW | | | |

RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GN, GQ, GW, ML, MR, NE, SN, TD, TG

US 2007087333 A1 20070419 US 2006-506418 20060818
PRIORITY APPLN. INFO.: EP 2004-75555 A 20040220
WO 2005-NL119 A1 20050218

AB The invention relates to a novel non-radioactive method to detect cytolytic activity against target cells expressing an antigenic antigen of choice. Cytotoxic T lymphocyte (CTL) activity provides a measure of the existence and magnitude of a cell-mediated cytotoxic response against a particular antigen. Specifically, the invention provides FATT (fluorescent-antigen-transfected target)-CTL assay, a kit and a nucleic acid for use in a method according to the invention. Cytotoxicity is quantified by assessing the elimination of viable cells expressing an antigen of interest associated with a fluorescent reporter mol., such as green fluorescent protein (GFP) expression assessing by flow cytometry. Thus, provided is a method for detecting cytolytic activity of cells or a substance against a population of target cells, comprising the steps of providing target cells with a first nucleic acid sequence encoding a reporter mol. and second nucleic acid sequence encoding an antigen of interest; co-culturing said target cells with a sample containing cells or a substance suspected of having cytolytic activity; and detecting the viability of target cells provided with the reporter mol. Demonstrated is use of FATT-CTL assay with HIV-1- and Influenza A virus-specific CTL and epitope variants.

IC ICM G01N033-569

CC 15-1 (Immunochromatography)

IT Antigen

RL: ARU (Analytical role, unclassified); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)

(Bacterial; FATT-CTL (fluorescent-antigen-transfected target -- cytotoxic T lymphocyte) assay, and kits to detect antigen-specific cytolytic activity for immunity assessing and drug screening)

IT Nucleic acids

RL: ANT (Analyte); BSU (Biological study, unclassified); ANST (Analytical study); BIOL (Biological study) (viability dye which stains; FATT-CTL (fluorescent-antigen-transfected target -- cytotoxic T lymphocyte) assay, and kits to detect antigen-specific cytolytic activity for immunity assessing and drug screening)

IT 157199-63-8, TO-PRO-3 iodide

RL: APG (Analytical reagent use); BSU (Biological study, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)

(FATT-CTL (fluorescent-antigen-transfected target -- cytotoxic T lymphocyte) assay, and kits to detect antigen-specific cytolytic activity for immunity assessing and drug screening)


IT 157199-63-8, TO-PRO-3 iodide

RL: APG (Analytical reagent use); BSU (Biological study, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)

(FATT-CTL (fluorescent-antigen-transfected target -- cytotoxic T lymphocyte) assay, and kits to detect antigen-specific cytolytic activity for immunity assessing and drug screening)

RN 157199-63-8 HCAPLUS

CN Quinolinium, 4-[3-(3-methyl-2(3H)-benzothiazolylidene)-1-propen-1-yl]-1-[3-(trimethylammonio)propyl]-, iodide (1:2) (CA INDEX NAME)



● 2 I -

REFERENCE COUNT:

7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L80 ANSWER 13 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 2005:601995 HCAPLUS Full-text
DOCUMENT NUMBER: 143:280037

TITLE: A self-contained fluorescent fiber optic DNA biosensor
AUTHOR(S): Wang, Xiaofeng; Krull, Ulrich J.
CORPORATE SOURCE: Chemical Sensors Group, Department of Chemical and Physical Sciences, University of Toronto, Mississauga, ON, L5L 1C6, Can.

SOURCE: Journal of Materials Chemistry (2005), 15(27-28), 2801-2809

PUBLISHER: CODEN: JMACEP; ISSN: 0959-9428
DOCUMENT TYPE: Royal Society of Chemistry
LANGUAGE: English

AB Single-stranded DNA (ssDNA) sequences can be used as probes to detect complementary targets, and represent useful anal. reagents for the detection and identification of bacteria, viruses and mutations. The hybridization process between probe sequences immobilized at a surface and complementary nucleic acid targets in a sample solution can, under optimal conditions, be complete in several minutes with a high degree of selectivity. Fluorescent dyes such as thiazole orange (TO) have been used extensively to quantify DNA by measuring the differential spectroscopic properties of free dye and the dye that assoc. with double-stranded DNA by intercalation. In an effort to develop a reagentless biosensor, TO has been covalently tethered by various poly(ether) strands at the 5' end of ssDNA probes, in a detection system where the oligonucleotide probes are immobilized onto the surfaces of fused silica optical fibers. Characterization of the surface immobilization has been completed using XPS. The biosensors provided changes in steady-state fluorescence intensity signals upon hybridization, that reached saturation in seconds to minutes, and were able to provide a quant. determination of hybridization at nanomolar detection limits. Aspects such as ionic strength, length of the tether that was used to attach TO to ssDNA, and the packing d. of the probe mols. were examined to determine the influence of these parameters on the thermodyn. and kinetic performance of the biosensor. In a

preliminary investigation of this application, the biosensor was used to detect PCR products from *Erwinia herbicola*.

CC 3-1 (Biochemical Genetics)

Section cross-reference(s): 9

IT 107091-89-4, Thiazole orange
RL: APG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)
(oligodeoxyribonucleotide conjugates; self-contained fluorescent fiber optic DNA biosensor)

IT 107091-89-4, Thiazole orange
RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)
(oligodeoxyribonucleotide conjugates; self-contained fluorescent fiber optic DNA biosensor)


RN 107091-89-4 HCAPLUS

CN Quinolinium, 1-methyl-4-[(3-methyl-2(3H)-benzothiazolylidene)methyl]-, 4-methylbenzenesulfonate (1:1) (CA INDEX NAME)

CM 1

CRN 24144-08-9


CMF C19 H17 N2 S



CM 2

CRN 16722-51-3

CMF C7 H7 O3 S



REFERENCE COUNT:

48 THERE ARE 48 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L80 ANSWER 14 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 2005:234346 HCAPLUS Full-text
DOCUMENT NUMBER: 142:409284

TITLE: $\gamma\delta$ T-lymphocyte cytotoxic activity against Mycobacterium bovis analyzed by flow cytometry

AUTHOR(S): Olin, Michael R.; Choi, K. Hwa; Lee, Jinhee; Molitor, Thomas W.

CORPORATE SOURCE: Clinical and Population Sciences, College of Veterinary Medicine, University of Minnesota, St. Paul, MN, 55108, USA

SOURCE: Journal of Immunological Methods (2005), 297(1-2), 1-11

CODEN: JIMMBG; **ISSN:** 0022-1759

PUBLISHER: Elsevier B.V.

DOCUMENT TYPE: English

LANGUAGE: English

AB $\gamma\delta$ T lymphocytes contain the unique capability of responding to pathogens in both an innate and acquired immune response. Previously, $\gamma\delta$ lymphocytes have been reported to respond to Mycobacteria tuberculosis determined by proliferation and IFN- γ production. Unlike $\alpha\beta$ lymphocytes, $\gamma\delta$ lymphocytes constitutively express a natural killer receptor providing $\gamma\delta$ lymphocytes the capability for innate cytolytic functions. A new cytolytic assay by flow cytometry was reported capable of determining natural killer activity using K562 cells as targets without the need for radioactive materials. The objectives of this study were to first apply the flow cytometer-based assay to assess $\gamma\delta$ lymphocytes natural killer activity following animal vaccination with Mycobacterium bovis Bacillus Calmette-Guérin (BCG). Secondly, to optimize the flow cytometer assay to detect antigen specific cytolytic activity to mycobacterium and to compare the cytolytic activity of $\gamma\delta$ lymphocytes to CD8 lymphocytes. $\gamma\delta$ lymphocytes increased in NK activity following animal vaccination with M. bovis BCG. Both innate and acquired antigen-specific cytolytic activity increased following incubation with M. bovis-infected monocytes. In conclusion, flow cytometric-based assay is a sensitive and reliable tool to determine cytolytic activity of $\gamma\delta$ T-lymphocytes against mycobacterium.

CC 15-1 (Immunochimistry)

ST T lymphocyte cytotoxicity Mycobacterium fluorescent dye flow cytometry

IT Infection

(bacterial; flow cytometric anal. of $\gamma\delta$ T-lymphocyte cytotoxic activity against Mycobacterium bovis)

IT T cell (lymphocyte)

(cytotoxic, TCR $\gamma\delta$; flow cytometric anal. of

$\gamma\delta$ T-lymphocyte cytotoxic activity against

Mycobacterium bovis)

IT Monocyte

(disease, infection; flow cytometric anal. of $\gamma\delta$

T-lymphocyte cytotoxic activity against Mycobacterium bovis)

IT Human

Immunofluorescence flow cytometry

Mycobacterium BCG

(flow cytometric anal. of $\gamma\delta$ T-lymphocyte cytotoxic

activity against Mycobacterium bovis)

IT Infection

(monocyte; flow cytometric anal. of $\gamma\delta$ T-lymphocyte

cytotoxic activity against Mycobacterium bovis)

IT 154214-55-8, PKH-26 157199-63-8, TO-PRO-3

RU: ANT (Analyte); ANST (Analytical study)

(flow cytometric anal. of $\gamma\delta$ T-lymphocyte cytotoxic

activity against Mycobacterium bovis)


IT 157199-63-8, TO-PRO-3

RL: ANT (Analyte); ANST (Analytical study)

(flow cytometric anal. of $\gamma\delta$ T-lymphocyte cytotoxic activity against Mycobacterium bovis)

RN 157199-63-8 HCAPLUS

CN Quinolium, 4-[(3-(3-methyl-2(3H)-benzothiazolylidene)-1-propen-1-yl)-1-(3-(trimethylammonio)propyl)]-, iodide (1:2) (CA INDEX NAME)



REFERENCE COUNT: 41

THERE ARE 41 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L80 ANSWER 15 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2004:780237 HCAPLUS Full-text

DOCUMENT NUMBER: 141:291842

TITLE: Dye compositions which provide enhanced differential fluorescence and light scatter characteristics

INVENTOR(S): Maples, John A.; Lopez, Lidice L.; Torke, Nancy

PATENT ASSIGNEE(S): Coulter International Corp., USA

SOURCE: U.S. Pat. Appl. Publ., 25 pp.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|---------------|--|----------|-----------------|----------|
| US 2004185447 | A1 | 20040923 | US 2003-392518 | 20030320 |
| US 6955872 | B2 | 20051018 | | |
| WO 2004085989 | A2 | 20041007 | WO 2004-US6889 | 20040305 |
| WO 2004085989 | A3 | 20041104 | | |
| W: | AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, GU, HK, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW | | | |
| RW: | BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, | | | |

ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GN, GW, ML, MR, NE, SN, TD, TG

EP 1604039 A2 20051214 EP 2004-718051 20040305
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK

JP 200520612 T 20060914 JP 2006-509203 20040305
PRIORITY APPLN. INFO.: US 2003-392518 A 20030320
WO 2004-US6889 W 20040305

AB A composition for enhancing differential staining of RNA, DNA and granules in a sample comprising cells contains a first fluorescent dye that can bind specific binding sites and non-specific binding sites in the sample. This first dye emits fluorescence at a first wavelength. The composition contains at least an addnl. component, which is a second non-intercalating dye in the composition that competes with said first dye for binding to the nonspecific binding sites, or a permeabilizing agent to enhance permeabilization of the dyes into the cells, or both. Reticulocytes were enumerated in blood samples by flow cytometry using a composition containing Acridine Orange as the primary dye, Hoechst 33258, and maltoside as sphering agent. The molar ratio of the second dye and the first dye is at least about 20:1.

IC ICM C12Q001-68
ICS C07H021-04
INCL 435006000; 536025320
CC 9-4 (Biochemical Methods)
ST dye enhanced differential fluorescence light scatter; RNA enhanced differential staining cell fluorescent dye; DNA enhanced differential staining cell fluorescent dye; granule enhanced differential staining cell fluorescent dye; reticulocyte flow cytometry Acridine Orange maltoside Hoechst 33258

IT Cyanine dyes
(BOBO; dye comps. which provide enhanced differential fluorescence and light scatter characteristics in staining RNA, DNA, and granules of cells)

IT Dyes
(SYTO; dye comps. which provide enhanced differential fluorescence and light scatter characteristics in staining RNA, DNA, and granules of cells)

IT Cyanine dyes
(TOYO; dye comps. which provide enhanced differential fluorescence and light scatter characteristics in staining RNA, DNA, and granules of cells)

IT Cyanine dyes
(YOYO; dye comps. which provide enhanced differential fluorescence and light scatter characteristics in staining RNA, DNA, and granules of cells)

IT Fluorescence
(autofluorescence; dye comps. which provide enhanced differential fluorescence and light scatter characteristics in staining RNA, DNA, and granules of cells)

IT Basophil
Blood analysis
Cell
Eosinophil
Erythrocyte
Flow cytometry
Fluorescent dyes
Human
Leukocyte
Lymphocyte
Mast cell

Monocyte
Neutrophil
Reticulocyte
Samples
Test kits
(dye comps. which provide enhanced differential fluorescence and light scatter characteristics in staining RNA, DNA, and granules of cells)

IT DNA
RNA
RL: BSU (Biological study, unclassified); DGN (Diagnostic use); BIOL (Biological study); USES (Uses)
(dye comps. which provide enhanced differential fluorescence and light scatter characteristics in staining RNA, DNA, and granules of cells)

IT Staining, biological
Stains, biological
(fluorescent; dye comps. which provide enhanced differential fluorescence and light scatter characteristics in staining RNA, DNA, and granules of cells)

IT Organelle
(granule; dye comps. which provide enhanced differential fluorescence and light scatter characteristics in staining RNA, DNA, and granules of cells)

IT Solvents
(in dye composition; dye comps. which provide enhanced differential fluorescence and light scatter characteristics in staining RNA, DNA, and granules of cells)

IT Diagnosis
(kit for; dye comps. which provide enhanced differential fluorescence and light scatter characteristics in staining RNA, DNA, and granules of cells)

IT Animal cell
(mammalian, blast; dye comps. which provide enhanced differential fluorescence and light scatter characteristics in staining RNA, DNA, and granules of cells)

IT Animal cell
(mammalian; dye comps. which provide enhanced differential fluorescence and light scatter characteristics in staining RNA, DNA, and granules of cells)

IT Staining, biological
(metachromasia; dye comps. which provide enhanced differential fluorescence and light scatter characteristics in staining RNA, DNA, and granules of cells)


IT Intercalating agents
(metachromatic dye as; dye comps. which provide enhanced differential fluorescence and light scatter characteristics in staining RNA, DNA, and granules of cells)

IT Dyes
(metachromatic; dye comps. which provide enhanced differential fluorescence and light scatter characteristics in staining RNA, DNA, and granules of cells)

IT Dyes
(nonintercalating, competing with first fluorescent dye for binding to nonspecific binding sites; dye comps. which provide enhanced differential fluorescence and light scatter characteristics in staining RNA, DNA, and granules of cells)

IT Erythrocyte
(nucleated; dye comps. which provide enhanced differential fluorescence and light scatter characteristics in staining

- IT Spheres
RNA, DNA, and granules of cells)
(permeabilizing agent as sphering agent; dye comps. which provide enhanced differential fluorescence and light scatter characteristics in staining RNA, DNA, and granules of cells)
- IT Biological transport
(permeation agent enhancing, in dye composition; dye comps. which provide enhanced differential fluorescence and light scatter characteristics in staining RNA, DNA, and granules of cells)
- IT Gallus domesticus
(red cells of; dye comps. which provide enhanced differential fluorescence and light scatter characteristics in staining RNA, DNA, and granules of cells)
- IT 65-61-2, Acridine Orange 83-89-6, Quinacrine 92-31-9, Toluidine Blue 92-32-0, Pyronin Y 135-57-9, Bisbenzamide 553-24-2, Neutral Red 1400-62-0, Orcein 2465-27-2, Auramine O 2465-29-4, Acridine Red 3056-93-7, Basic Orange 21 6493-05-6, Pentoxiphylline 17372-87-1, Eosin Y 18472-87-2, Phloxine B 23491-45-4, Hoechst 33258 23491-52-3, Hoechst 33342 23555-00-2, Hoechst 34580 25535-16-4, Propidium iodide 47165-04-8, DAPI 48198-86-3D, derivs. 64431-93-2 75168-11-5, Acridine Orange 10-nonyl bromide 76433-29-9, LDS 751 107091-89-4, Thiazole Orange 157199-63-8, TO-PRO-3
RL: ARG (Analytical reagent use); BSU (Biological study, unclassified); BUU (Biological use, unclassified); DGN (Diagnostic use); ANST (Analytical study); BGIOL (Biological study); USES (Uses)
(as dye; dye comps. which provide enhanced differential fluorescence and light scatter characteristics in staining RNA, DNA, and granules of cells)
- IT 14933-08-5, n-Dodecyl-N,N-dimethyl-3-ammonio-1-propanesulfonate 69227-93-6, n-Dodecyl-β-D-maltoside
RL: ARG (Analytical reagent use); BSU (Biological study, unclassified); BUU (Biological use, unclassified); DGN (Diagnostic use); ANST (Analytical study); BGIOL (Biological study); USES (Uses)
(as sphering agent in dye composition; dye comps. which provide enhanced differential fluorescence and light scatter characteristics in staining RNA, DNA, and granules of cells)
- IT 76433-29-9, LDS 751 107091-89-4, Thiazole Orange 157199-63-8, TO-PRO-3
RL: ARG (Analytical reagent use); BSU (Biological study, unclassified); BUU (Biological use, unclassified); DGN (Diagnostic use); ANST (Analytical study); BGIOL (Biological study); USES (Uses)
(as dye; dye comps. which provide enhanced differential fluorescence and light scatter characteristics in staining RNA, DNA, and granules of cells)
- RN 76433-29-9 HCAPLUS
CN Benzo[thiazolium, 2-{4-[4-(dimethylamino)phenyl]-1,3-butadien-1-yl]-3-ethyl-, perchlorate (1:1) (CA INDEX NAME)
- CM 1
CRN 76433-28-8
CMF C21 H21 N2 S




CM 2
CRN 14797-73-0
CMF C1 O4




RN 107091-89-4 HCAPLUS
CN Quinolinium, 1-methyl-4-((3-methyl-2-(3H)-benzothiazolylidene)methyl)-, 4-methylbenzenesulfonate (1:1) (CA INDEX NAME)


CM 1
CRN 24144-08-9
CMF C19 H17 N2 S



CM 2
CRN 16722-51-3
CMF C7 H7 O3 S



RN 157199-63-8 HCAPLUS
CN Quinolium, 4-[(3-methyl-2(3H)-benzothiazolylidene)-1-propen-1-yl]-1-(3-(trimethylammonio)propyl)-, iodide (1:2) (CA INDEX NAME)



● 2 I⁻

REFERENCE COUNT: 19 THERE ARE 19 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L80 ANSWER 16 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 2004:589685 HCAPLUS Full-text
DOCUMENT NUMBER: 141:118285
TITLE: Use of sensor arrays containing hairpin probes for detecting nucleic acids of pathogens
INVENTOR(S): Miller, Benjamin L.; Krauss, Todd D.; Du, Hui; Crnkovich, Nicole; Strohsahl, Christopher M.
PATENT ASSIGNEE(S): University of Rochester, USA
SOURCE: PCT Int. Appl., 73 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|-----------------------|------|--|-----------------|------------|
| WO 2004061127 | A2 | 20040722 | WO 2004-US93 | 20040102 |
| WO 2004061127 | A3 | 20050630 | | |
| | W: | AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, W: | | |
| CA 2511874 | A1 | 20040722 | CA 2004-2511874 | 20040102 |
| US 2007059693 | A1 | 20070315 | US 2005-541044 | 20050624 |
| PRIORITY APPL. INFO.: | | | US 2003-43780P | P 20030102 |
| | | | WO 2004-US93 | W 20040102 |

AB The present invention provides use of sensor arrays containing hairpin probes for detecting nucleic acids of pathogens. Various nucleic acid probes,

methods of making the sensor chip, biol. sensor devices that contain the sensor chip, and their methods of use are also disclosed.

IC ICM CL2Q
CC 3-1 (Biochemical Genetics)
IT Section cross-reference(s): 9, 10, 14
Actinobacter calcoaceticus
Actinobacillus
Adenoviridae
Aeromonas hydrophila
Arbovirus
Arizona hinhawii
Ateline herpesvirus
Avalanche photodiodes
Avian leukosis virus
Bacillus anthracis
Bartonella
Biomarkers
Biosensors
Blastomyces dermatitidis
Bordetella
Borrelia
Bovine leukemia virus
Bovine papillomavirus
Brucella
Cache Valley virus
Campylobacter
Charge coupled devices
Chlamydia
Clostridium
Coccidioides immitis
Coronavirus
Corynebacterium
Cryptococcus neoformans
Cytomegalovirus
DNA microarray technology
Dengue virus
Dermatophilus congolensis
Disease, animal
Ebola virus
Edwardsiella tarda
Encephalomyocarditis virus
Entamoeba histolytica
Erysipelothrix rhusiopathiae
Escherichia coli
Eubacteria
Feline leukemia virus
Feline sarcoma virus
Flanders virus
Fluorometry
Fowl adenovirus 1
Francisella tularensis
Fungi
Pseudomonas aeruginosa
Pseudomonas necrophorum
Gallid herpesvirus
Genetic polymorphism
Haemophilus
Hart Park virus
Hepatitis virus
Herpes virus B
Herpesviridae

Histoplasma
Human
Human coxsackievirus A
Human coxsackievirus B
Human echovirus
Human herpesvirus 3
Human herpesvirus 4
Human parainfluenza virus
Human poliovirus
Influenza virus
Klebsiella
Langat virus
Lasers
Lassa virus
Legionella pneumophila
Leishmania
Leptospira interrogans
Listeria
Lymphocytic choriomeningitis virus
Marburg virus
Mason-Pfizer monkey virus
Measles virus
Monkeypox virus
Moraxella
Mouse mammary tumor virus
Mumps virus
Murine leukemia virus
Murine sarcoma virus
Mycobacterium
Mycobacterium avium
Mycoplasma
Naegleria gruberi
Neisseria
Nocardia
Nucleic acid hybridization
Paracoccidioides brasiliensis
Parasite
Pasteurella
Pathogen
Photodiodes
Photomultipliers
Pneumocystis carinii
Polyomavirus
Poxviridae
Pseudomonas
Pseudonocardia autotrophica
Rabbit fibroma virus
Rabies virus
Rat leukemia virus
Reoviridae
Respiratory syncytial virus
Rhinovirus
Rhodococcus
Rous sarcoma virus
Rubella virus
Saimiriine herpesvirus
Salmonella
Schistosoma mansoni
Shigella
Shope papilloma virus

Simian virus 40
Sindbis virus
Staphylococcus aureus
Streptococcus moniliformis
Streptococcus
Tensaw virus
Tick-borne encephalitis virus
Toxocara canis
Toxoplasma gondii
Treponema
Trichinella spiralis
Trypanosoma cruzi
Turlock virus
Vaccinia virus
Venezuelan equine encephalitis virus
Vesicular stomatitis virus
Vibrio
Virus
Woolly monkey sarcoma virus
Yaba monkey tumor virus
Yellow fever virus
Yersinia

(use of sensor arrays containing hairpin probes for detecting nucleic acids of pathogens)

IT 81-88-9D, RhodamineB, probe conjugate 92-32-0D, Pyronin Y, probe conjugate 1461-15-0D, Calcein, probe conjugate 7385-67-3D, Nile red, probe conjugate 13558-31-1D, Rhodamine 110, probe conjugate 27072-45-3D, Fitt, probe conjugate 41085-99-8D, DiI, probe conjugate 62669-70-9D, Rhodamine 123, probe conjugate 76823-03-5D, 5-Carboxyfluorescein, probe conjugate 82354-19-6D, Texas red, probe conjugate 82855-40-1D, Joe, probe conjugate 99752-92-8D, Rhodamine red, probe conjugate 107347-53-5D, Tric, probe conjugate 120718-39-0D, Rox, probe conjugate 120718-52-7D, Tamra, probe conjugate 127274-91-3D, DiD, probe conjugate 138067-54-6D, Calcium crimson, probe conjugate 138067-55-7D, Calcium green, probe conjugate 138067-56-8D, Calcium orange, probe conjugate 143413-84-7D, TOPO-1, probe conjugate 143413-85-8D, Yoyo-1, probe conjugate 150173-72-1D, Bodipy 558/568, probe conjugate 150173-89-0D, Bodipy 564/570, probe conjugate 152068-09-2D, Yo-pro-1, probe conjugate 156312-20-8D, Yoyo-3, probe conjugate 157199-59-2D, TO-PRO-1, probe conjugate 157199-62-7D, Yo-pro-3, probe conjugate 157199-63-8D, To-pro-3, probe conjugate 166196-17-4D, Toto-3, probe conjugate 170516-41-3D, Magnesium green, probe conjugate 172777-84-3D, Cys 5, probe conjugate 187089-10-7D, BODIPY 530/550, probe conjugate 189200-71-3D, Rhodamine green, probe conjugate 189767-45-1D, Cy3.5, probe conjugate 189767-52-0D, FluorX, probe conjugate 195136-58-4D, Oregon green 488, probe conjugate 220751-06-4D, Ribogreen, probe conjugate 247145-23-9D, Alexa 546, probe conjugate 247145-86-4D, Alexa 594, probe conjugate 915013-10-4D, Rhodamine phalloidin, probe conjugate

RL: ARG (Analytical reagent use); DGN (Diagnostic use)
; ANST (Analytical study); BIOL (Biological study);
USES (Uses)

(use of sensor arrays containing hairpin probes for detecting nucleic acids of pathogens)

IT 157199-63-8D, To-pro-3, probe conjugate 166196-17-4D, Toto-3, probe conjugate
RL: ARG (Analytical reagent use); DGN (Diagnostic use)
; ANST (Analytical study); BIOL (Biological study);
USES (Uses)


(use of sensor arrays containing hairpin probes for detecting nucleic acids

of pathogens)

RN 157199-63-8 HCAPLUS

CN Quinolium, 4-[(3-methyl-2(3H)-benzothiazolylidene)-1-propen-1-yl]-1-[(3-trimethylammonio)propyl]-, iodide (1:2) (CA INDEX NAME)


(CH₂)₃-N⁺-Me₃



● 2 I-


RN 166196-17-4 HCAPLUS

CN Quinolium, 1,1'-[1,3-propanediylbis[(dimethyliminio)-3,1-propanediyl]bis[4-(3-methyl-2(3H)-benzothiazolylidene)-1-propen-1-yl]-, iodide (1:4) (CA INDEX NAME)



PAGE 1-A

PAGE 2-A



L80 ANSWER 17 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2004:392699 HCAPLUS Full-text
 DOCUMENT NUMBER: 140:371469
 TITLE:
 A method for assessment of particles
 INVENTOR(S): Larsen, Rasmus Dines; Hansen, Frans Ejner Ravn
 PATENT ASSIGNEE(S): Chemometec A/s, Den.
 SOURCE: PCT Int. Appl., 70 pp.
 CODEN: PIXXD2

DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|---|------|----------|-----------------|----------|
| WO 2004040314 | A1 | 20040513 | WO 2003-DK743 | 20031031 |
| W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KZ, LC, LK, LR, LS, LT, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SN, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW | | | | |
| RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GN, GQ, GW, ML, MR, NE, SN, TD, TG | | | | |
| AU 2003275940 | A1 | 20040525 | AU 2003-275940 | 20031031 |
| EP 1558934 | A1 | 20050803 | EP 2003-809704 | 20031031 |
| R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK | | | | |
| JP 2006504937 | T | 20060209 | JP 2004-547451 | 20031031 |
| US 2006063146 | A1 | 20060323 | US 2005-533324 | 20050812 |
| PRIORITY APPLN. INFO.: DK 2002-1653 A 20021031 | | | | |
| WO 2003-DK743 W 20031031 | | | | |

AB The invention relates to imaging methods for assessing quality or quantity parameters of particles in a sample, wherein the particles contain less than 106 analyte detectable positions. The method comprises (1) mixing the sample with a targeting species capable of binding an analyte position and a labeling agent, (2) arranging the sample in an exposing domain, allowing

electromagnetic signals from the sample to pass to the exterior, (3) exposing a representation of said signals onto an array of detection elements, wherein the representation is subject to a linear enlargement, so that the ratio of the image of a linear dimension on the array of detection elements to the original linear dimension in the exposing domain is smaller than 20:1, (4) detecting the representation as intensities by said detection elements, (5) processing the intensities to identify the particles, and (6) obtaining the quality or quantity parameter.

IC ICM GOIN033-58
CC 9-4 (Biochemical Methods)

IT Blood cell

Cell
Eubacteria

Human
Nucleic acid hybridization
Staining, biological

Virus

cDNA sequences

(method for assessment of particles)

IT 65-61-2, Acridine orange 2321-07-5, Fluorescein 7240-37-1,
7-Aminoactinomycin d 25535-16-4, Propidium iodide 27072-45-3, FITC
47165-04-8, DAPI 64457-77-8, Ethidium iodide 107091-89-4,

Thiazole orange 146368-14-1, Cy5 146368-16-3, Cy3 172777-84-3, Cy5.5

RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)

(method for assessment of particles)

IT 107091-89-4, Thiazole orange

RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)

(method for assessment of particles)


RN 107091-89-4 HCAPLUS

CN Quinolium, 1-methyl-4-[(3-methyl-2(3H)-benzothiazolylidene)methyl]-,
4-methylbenzenesulfonate (1:1) (CA INDEX NAME)

CM 1

CRN 24144-08-9


CMF C19 H17 N2 S



CM 2

CRN 16722-51-3

CMF C7 H7 O3 S



REFERENCE COUNT: 5

THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L80 ANSWER 18 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 2003:757324 HCAPLUS Full-text
DOCUMENT NUMBER: 139:272000

TITLE:
Method for modifying transcription and/or translation
in an organism by heteropolymetric probes and duplex,
triplex or quadruplex hybridization for therapeutic,
prophylactic and/or analytic uses

INVENTOR(S):

Erikson, Glen H.

PATENT ASSIGNEE(S):

Ingeneus Corporation, Barbados

U.S. Pat. Appl. Publ., 54 pp., Cont.-in-part of U.S.

SOURCE:

Ser. No. 909,496.

CODEN: USXXCO

DOCUMENT TYPE:

Patent

LANGUAGE:


English

FAMILY ACC. NUM. COUNT: 15

PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|---|------|----------|-----------------|-------------|
| US 2003181412 | A1 | 20030925 | US 2003-438151 | 20030514 |
| US 6403313 | B1 | 20020611 | US 1999-468679 | 19991221 |
| US 6420115 | B1 | 20020716 | US 2000-613263 | 20000710 |
| US 6903300 | B1 | 20050531 | US 2000-664827 | 20000919 |
| US 2002031775 | A1 | 20020314 | US 2001-909496 | 20010720 |
| US 6656692 | B2 | 20031202 | | |
| WO 2004100636 | A2 | 20041125 | WO 2003-1B5624 | 20030924 |
| WO 2004100636 | A3 | 20050317 | | |
| W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, GU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TD, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW | | | | |
| RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GW, ML, MR, NE, SN, TD, TG | | | | |
| AU 2003282322 | A1 | 20041203 | AU 2003-282322 | 20030924 |
| ZA 2005009517 | A | 20060726 | ZA 2005-9517 | 20051124 |
| PRIORITY APPLN. INFO.: | | | | |
| | | | US 1999-468679 | A2 19991221 |
| | | | US 2000-613263 | A2 20000710 |
| | | | US 2000-664827 | A2 20000919 |
| | | | US 2001-909496 | A2 20010720 |
| | | | US 2003-438151 | A 20030514 |
| | | | WO 2003-1B5624 | W 20030924 |

A5 The invention relates to a method for modifying gene expression for therapeutic and/or prophylactic purposes, and more particularly to such a method wherein duplex, triplex and/or quadruplex complexes are formed by specific binding between single-stranded or double-stranded nucleobase-



L80 ANSWER 19 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2003:605001 HCAPLUS Full-text

DOCUMENT NUMBER: 140:283673

TITLE: Optimization of three- and four-color multiparameter DNA analysis in lymphoma specimens

AUTHOR(S): Plander, M.; Brockhoff, G.; Barlage, S.; Schwarz, S.; Rothe, G.; Kneuchel, R.

CORPORATE SOURCE: Department of Hematology, University Teaching Hospital of Vas County, Szombathely, Hung.

SOURCE: Cytometry, Part A (2003), 54A(1), 66-74

CODEN: CPAVAV

PUBLISHER: Wiley-Liss, Inc.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Background: Simultaneous anal. of DNA and immunophenotype of lymphoma cells by flow cytometry allows the calcn. of the proliferative activity and aneuploidy in even a small lymphoma population. Unfavorable DNA binding characteristics or spectral features of DNA dyes impair the accuracy of multiparameter DNA anal. and limit their clin. application. We describe here a reliable and reproducible application of both three- and four-color multiparameter DNA anal. Methods: After immunostaining of fresh samples of peripheral blood, bone marrow and single cell suspensions of lymph nodes from healthy and lymphoma patients, a methanol fixation for TO-PRO-3 and DRAQ5 staining was tested. Results: The red-excitable TO-PRO-3 on a FACScalibur is limited to two-color antigen staining including fluorescein-isothiocyanate and phycoerythrin-labeled monoclonal antibodies due to its broad excitation spectrum. Although DRAQ5 is only applicable to flow cytometers equipped with a single argon laser emitting 488-nm light, its emission spectrum can be easily separated from the FITC, PE, and PE/Texas-Red emissions. DRAQ5 showed almost identical stoichiometric DNA binding characteristics as propidium iodide. Coefficient of variation produced by DRAQ5 staining is in the range of 3.5 and is adequate for detecting aneuploid and near-diploid cells. Conclusions: These advantageous features of DRAQ5 make it a reliable candidate for multiparameter clin. studies.

CC 9-5 (Biochemical Methods)

Section cross-reference(s): 6, 13, 14

ST DNA fluorescence staining flow cytometry lymphoma diagnosis

IT Staining, biological

Stains, biological

(fluorescent; flow cytometry three- and four-color multiparameter DNA anal. in lymphoma specimens)

IT 25535-16-4, Propidium iodide 27072-45-3, FITC 82354-19-6, Texas Red

157199-63-8, To-Pro 3 254098-36-7, DRAQ5 422551-33-5, PerCP

RL: ARU (Analytical role, unclassified); BUU (Biological

use, unclassified); ANST (Analytical study); BIOL

(Biological study); USES (Uses)

(flow cytometry three- and four-color multiparameter DNA anal. in

lymphoma specimens)

IT 157199-63-8, To-Pro 3

RL: ARU (Analytical role, unclassified); BUU (Biological

use, unclassified); ANST (Analytical study); BIOL

(Biological study); USES (Uses)


(flow cytometry three- and four-color multiparameter DNA anal. in

lymphoma specimens)

RN 157199-63-8 HCAPLUS

CN Quinolinium, 4-[3-(3-methyl-2(3H)-benzothiazolylidene)-1-propen-1-yl]-1-[3-

(trimethylammonio)propyl]-, iodide (1:2) (CA INDEX NAME)



REFERENCE COUNT: 33

THERE ARE 33 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L80 ANSWER 20 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2003:499067 HCAPLUS Full-text

DOCUMENT NUMBER: 140:267094

TITLE: Interaction of cyanine dyes with nucleic acids: XXXI. Using of polymethine cyanine dyes for the visualization of DNA in agarose gels

Matselyukh, B. P.; Yarmoluk, S. M.; Matselyukh, A. B.; Kovalska, V. B.; Kocheshev, I. O.; Kryvorotenko, D. V.; Lukashov, S. S.

CORPORATE SOURCE: Institute of Microbiology and Virology of the National Academy of Sciences of Ukraine, Kiev, 03143, Ukraine

SOURCE: Journal of Biochemical and Biophysical Methods (2003), 57(1), 35-43

PUBLISHER: Elsevier Science Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Fifteen polymethine cyanine dyes were studied as fluorescent stains for DNA in electrophoretic gels. Among studied cyanines, two dyes CPent V and CCyan 2-O most effectively visualized covalently closed and linear double-stranded DNA mols. in gels under standard conditions using UV-illumination, green filter and black-and-white photo film. Ethidium bromide was 1.2-1.6 times more effective as compared to cyanine dyes in staining of DNA in the concentration range of 8-16 ng, while studied cyanines were more sensitive to DNA quantity above 50 ng.

CC 9-16 (Biochemical Methods)

IT Fluorescent dyes

IT (cyanine); polymethine cyanine dyes for fluorescent staining and visualization of DNA in agarose gels)

IT DNA

RL: ANT (Analyte); BSU (Biological study, unclassified); ANST (Analytical study); BIOL (Biological study)

(double-stranded; polymethine cyanine dyes for fluorescent staining and visualization of DNA in agarose gels)

IT Cyanine dyes

Staining, biological

Stains, biological

(fluorescent; polymethine cyanine dyes for fluorescent staining and visualization of DNA in agarose gels)

IT Fluorometry

Gel electrophoresis

Molecular association

(polymethine cyanine dyes for fluorescent staining and visualization of DNA in agarose gels)

IT DNA

RL: ANT (Analyte); BSU (Biological study, unclassified); ANST (Analytical study); BIOL (Biological study)

(polymethine cyanine dyes for fluorescent staining and visualization of DNA in agarose gels)

IT

1239-45-8, Ethidium bromide 7423-31-6, Stains-All

106396-46-7 107091-89-4 287966-83-0 33380-50-0

33380-52-2 33380-56-6 33380-61-3 340157-38-2 349081-16-9

351526-14-2 485403-76-7 497220-85-6 569361-79-1 674784-62-4

674784-65-7

RL: ARG (Analytical reagent use); BSU (Biological study, unclassified); ANST (Analytical study); BIOL

(Biological study); USES (Uses)

(polymethine cyanine dyes for fluorescent staining and visualization of DNA in agarose gels)

IT

107091-89-4

RL: ARG (Analytical reagent use); BSU (Biological study, unclassified); ANST (Analytical study); BIOL

(Biological study); USES (Uses)

(polymethine cyanine dyes for fluorescent staining and visualization of DNA in agarose gels)

107091-89-4 HCAPLUS


RN Quinolium, 1-methyl-4-[(3-methyl-2(3H)-benzothiazolylidene)methyl].

CN 4-methylbenzenesulfonate (1:1) (CA INDEX NAME)

CM 1

CRN 24144-08-9


CMF C19 H17 N2 S



CM 2

CRN 16722-51-3

CMF C7 H7 O3 S



REFERENCE COUNT: 16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L80 ANSWER 21 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2002-450245 HCAPLUS Full-text

DOCUMENT NUMBER: 137:30238

TITLE: Immunoassay based on DNA replication using labeled primer

INVENTOR(S): McNally, Alan J.; Wu, Robert S.; Li, Zhuyin

PATENT ASSIGNEE(S): USA

SOURCE: U.S. Pat. Appl. Publ., 26 pp.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO. KIND DATE APPLICATION NO. DATE

US 2002072053 A1 20020613 US 2000-733565 20001208

PRIORITY APPLN. INFO.: US 2000-733565 20001208

AB The invention concerns an immunoassay method based upon inhibition of a DNA

polymerase enzyme accomplished by linking a ligand of the analyte to a primer

through a covalent bond. The interaction between the primer-bound ligand and

a receptor specific for the ligand inhibits the DNA polymerase enzyme from

generating double stranded DNA. The degree of inhibition of double stranded

DNA synthesis is inversely proportional to the concentration of analyte in the

sample. The analyte is determined by measuring the formation of double

stranded DNA, e.g., by a fluorescence DNA intercalation technique.

IC ICM C12Q001-68

INCL 435006000

CC 9-10 (Biochemical Methods)

bacteria, and a diluent for bacterial stain

IT Functional groups
(benzyl group; method of staining, detecting and counting bacteria, and a diluent for bacterial stain)

IT Surfactants
(cationic; method of staining, detecting and counting bacteria, and a diluent for bacterial stain)

IT Anions
(comps. containing; method of staining, detecting and counting bacteria, and a diluent for bacterial stain)

IT Measuring apparatus
(cytometers, flow; method of staining, detecting and counting bacteria, and a diluent for bacterial stain)

IT Halogens
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)
(ions; method of staining, detecting and counting bacteria, and a diluent for bacterial stain)

IT Acyl groups
Blood analysis
Body fluid
Buffers
Cerebrospinal fluid
Chemical formula
Cyanine dyes
Dilution
Dyes
Eubacteria
Flow cytometry
Fluorometry
Length
Light
Light scattering
Optical reflection
Particles
Radiation
Reducing agents
Samples
Solutions
Staining, biological
Stains, biological
Urine analysis
PH
(method of staining, detecting and counting bacteria, and a diluent for bacterial stain)

IT Acids, biological studies
RL: BSU (Biological study, unclassified); BIOL (Biological study)
(method of staining, detecting and counting bacteria, and a diluent for bacterial stain)

IT Nitrates, biological studies
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)
(method of staining, detecting and counting bacteria, and a diluent for bacterial stain)

IT Quaternary ammonium compounds, biological studies
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)
(method of staining, detecting and counting bacteria, and a diluent for bacterial stain)

IT Salts, biological studies
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)
(method of staining, detecting and counting bacteria, and a diluent for bacterial stain)

IT Sulfates, biological studies
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)
(method of staining, detecting and counting bacteria, and a diluent for bacterial stain)

IT 14797-65-0, Nitrite ion, biological studies
RL: BSU (Biological study, unclassified); RCT (Reactant); BIOL (Biological study); RCT (Reactant or reagent)
(method of staining, detecting and counting bacteria, and a diluent for bacterial stain)

IT 50-21-5, Lactic acid, biological studies 56-40-6, Glycine, biological studies 60-32-2, L-Alanine 110-15-6, Succinic acid, biological studies 107-95-9, β -Alanine 110-15-6, Succinic acid, biological studies 110-17-8, Fumaric acid, biological studies 877-24-7, Potassium hydrogen phthalate 1119-97-7, Tetradecyl trimethylammonium bromide 1310-71-2, Sodium hydroxide (NaOH), biological studies 1333-74-0D, Hydrogen, compds. containing 6899-10-1 7440-44-0D, Carbon, compds. containing 7558-79-4, Disodium hydrogen phosphate 7647-01-0, Hydrochloric acid, biological studies 7704-34-9D, Sulfur, compds. containing 7778-77-0, Potassium dihydrogen phosphate 7782-44-7D, Oxygen, compds. containing 10182-91-9 10182-92-0 15053-09-5 15461-40-2 76433-27-7 76433-29-9 150749-57-5 157189-63-8 166196-17-4 189148-50-3 335080-22-3 361544-71-0 361544-72-1

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)
(method of staining, detecting and counting bacteria, and a diluent for bacterial stain)

IT 50-81-7, Ascorbic acid, biological studies 52-90-4, Cysteine, biological studies 56-84-8, Aspartic acid, biological studies 56-85-9, Glutamine, biological studies 56-86-0, Glutamic acid, biological studies 57-13-6, Urea, biological studies 60-24-2, Mercaptoethanol 63-68-3, Methionine, biological studies 63-74-1, Sulfanilamide 68-11-1, Mercaptoacetic acid, biological studies 70-18-8, Glutathione, biological studies 70-47-3, Asparagine, biological studies 74-89-5, Aminomethane, biological studies 89-65-6, Isoascorbic acid 107-35-7 107-96-0, 3-Mercaptopropionic acid 108-98-5, Thiophenol, biological studies 121-57-3, Sulfanilic acid 5329-14-6, Sulfamic acid 6303-21-5, Phosphonic acid 7782-99-2, Sulfurous acid, biological studies 7803-49-8, Hydroxylamine, biological studies 7803-49-8D, Hydroxylamine, salts 13881-91-9, Aminomethanesulfonic acid 33669-61-3, Pyrosulfurous acid

RL: BUU (Biological use, unclassified); RCT (Reactant); BIOL (Biological study); RCT (Reactant or reagent); USES (Uses)
(method of staining, detecting and counting bacteria, and a diluent for bacterial stain)

IT 24147-35-2, Thiazole orange
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)
(thiazole orange; method of staining, detecting and counting


August 23, 2007

10/803,667

bacteria, and a diluent for bacterial strain

IT 76433-27-7 76433-29-9 150749-57-8
 157199-63-8 166196-17-4 189148-50-3
 335080-22-3 361544-71-0 361544-72-1
 RL: BUU (Biological use, unclassified); BIOL (Biological
 study); USES (Uses)
 (method of staining, detecting and counting bacteria
 , and a diluent for bacterial strain)
 RN 76433-27-7 HCAPLUS
 CN 3H-Indolium, 2-[4-[4-(dimethylamino)phenyl]-1,3-butadienyl]-1,3,3-
 trimethyl-, perchlorate (9CI) (CA INDEX NAME)

CM 1
 CRN 76433-26-6
 CMF C23 H27 N2



CM 2
 CRN 14797-73-0
 CMF Cl O4



RN 76433-29-9 HCAPLUS
 CN Benzothiazolium, 2-[4-[4-(dimethylamino)phenyl]-1,3-butadien-1-yl]-3-ethyl-
 , perchlorate (1:1) (CA INDEX NAME)

CM 1
 CRN 76433-28-8
 CMF C21 H23 N2 S



55


August 23, 2007

10/803,667


CM 2
 CRN 14797-73-0
 CMF Cl O4



RN 150749-57-8 HCAPLUS
 CN Benzothiazolium, 3-[3-(trimethylammonio)propyl]-2-[5-[3-(3-
 (trimethylammonio)propyl]-2(3H)-benzothiazolylidene]-1,3-pentadienyl]-,
 tribromide (9CI) (CA INDEX NAME)



RN 157199-63-8 HCAPLUS
 CN Quinolium, 4-[3-(3-methyl-2(3H)-benzothiazolylidene)-1-propen-1-yl]-1-[3-
 (trimethylammonio)propyl]-, iodide (1:2) (CA INDEX NAME)




56

10/803,667


August 23, 2007

RN 166196-17-4 HCAPLUS
CN Quinolinium, 1,1'-(1,3-propanediylbis[(dimethyliminio)-3,1'-propanediyl]bis[4-(3-(3-methyl-2(3H)-benzothiazolylidene)-1-propen-1-yl]-, iodide (1:4) (CA INDEX NAME)

PAGE 1-A



PAGE 2-A



RN 189148-50-3 HCAPLUS
CN Quinolinium, 1-(2-hydroxyethyl)-4-(3-(3-methyl-2(3H)-benzothiazolylidene)-1-propenyl-, tetrafluoroborate (1-) (9CI) (CA INDEX NAME)


CM 1

57

10/803,667

August 23, 2007

CRN 189148-49-0
CMF C22 H21 N2 O S



CM 2

CRN 14874-70-5
CMF B F4
CCI CCS



RN 335080-22-3 HCAPLUS


CN Benzenesulfonic acid, 4-[4-[5-(1,3-dibutylhexahydro-4,6-dioxo-2-thioxo-5-pyrimidinyl)-2,4-pentadienylidene]-4,5-dihydro-3-methyl-5-oxo-1H-pyrazol-1-yl]-, compd. with N,N-diethylethanamine (1:2) (9CI) (CA INDEX NAME)

CM 1


CRN 118702-42-4
CMF C27 H32 N4 O6 S2

58


PAGE 1-A




PAGE 1-A



PAGE 1-B



PAGE 2-A




CM 2

CRN 121-44-8
CMF C6 H15 N


Et
Et-N-Et

RN 361544-71-0 HCAPLUS
CN Borate(1-), difluoro[2,3,5,6-tetrafluoro-4-sulphophenyl
6-[[[4-[2-[5-(2-thienyl)-2H-pyrrol-2-ylidene-KN]methyl]-1H-
pyrrol-2-yl-KN]ethenyl]phenoxy]acetyl]amino]hexanoato(2-)]-, sodium,
(T-4) - (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 1-B



IT 24147-36-2, Thiazole orange
 RL: BUU (Biological use, unclassified); BIOL (Biological study); USSS (Uses)
 (thiazole orange; method of staining, detecting and counting bacteria, and a diluent for bacterial stain)

RN 24147-36-2 HCAPLUS

CN Quinolinium, 1-methyl-4-[(3-methyl-2(3H)-benzothiazolylidene)methyl]-, iodide (1:1) (CA INDEX NAME)




● I⁻

L80 ANSWER 23 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2001:764185 HCAPLUS Full-text
 DOCUMENT NUMBER: 136:289517
 TITLE: PNA-based light-up probes for real-time detection of sequence-specific PCR products
 AUTHOR(S): Wolffs, Petra; Knutsson, Rickard; Sjoback, Robert; Radstrom, Peter
 CORPORATE SOURCE: Lund University, Lund, Swed.
 SOURCE: BioTechniques (2001), 31(4), 766, 769-771
 CODEN: BTNQDQ; ISSN: 0736-6205
 PUBLISHER: Eaton Publishing Co.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB The aim of this study was to introduce the use of a peptide nucleic acid (PNA)-thiazole orange conjugate for real-time monitoring of PCR. When the so-called light-up probes hybridize sequence-specifically to the PCR product, an increase in the fluorescent signal is obtained. It was found that the light-up probe can quant. measure the amount of DNA or intact bacterial cells in the reaction mixture, without interfering with the PCR amplification. A linear detection range of at least 4 log units was obtained without optimization of the system. The detection limit of this light-up assay per reaction mixture was 0.4 pg genomic *Yersinia enterocolitica* DNA.
 CC 3-1 (Biochemical Genetics)
 IT DNA

RL: ANT (Analyte); ANST (Analytical study)
 (crude bacterial extract or purified; PNA-based light-up probes for real-time detection of sequence-specific PCR products)
 IT 24147-36-2D, Thiazole orange, conjugates with peptide nucleic acid
 RL: ARG (Analytical reagent use); ANST (Analytical

study); USSS (Uses)
 (thiazole orange; PNA-based light-up probes for real-time detection of sequence-specific PCR products)
 IT 24147-36-2D, Thiazole orange, conjugates with peptide nucleic acid
 RL: ARG (Analytical reagent use); ANST (Analytical study); USSS (Uses)
 (thiazole orange; PNA-based light-up probes for real-time detection of sequence-specific PCR products)
 RN 24147-36-2 HCAPLUS
 CN Quinolinium, 1-methyl-4-[(3-methyl-2(3H)-benzothiazolylidene)methyl]-, iodide (1:1) (CA INDEX NAME)



● I⁻

REFERENCE COUNT: 20 THERE ARE 20 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L80 ANSWER 24 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2001:702413 HCAPLUS Full-text
 DOCUMENT NUMBER: 135:254110
 TITLE: Method for staining and detecting bacteria
 INVENTOR(S): Inoue, Junya; Ikeuchi, Yoshiro; Kawashima, Yasuyuki
 PATENT ASSIGNEE(S): Sysmex Co., Ltd., Japan
 SOURCE: Jpn. Kokai Tokkyo Koho, 11 pp.
 CODEN: JKXXAF

DOCUMENT TYPE: Patent
 LANGUAGE: Japanese
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|---------------|------|----------|-----------------|----------|
| JP 2001258590 | A | 20010925 | JP 2000-80998 | 20000322 |
| JP 3837006 | B2 | 20061025 | | |
| EP 1136563 | A2 | 20010926 | EP 2001-201027 | 20010320 |
| EP 1136563 | A3 | 20040121 | | |
| EP 1136563 | B1 | 20060607 | | |

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, CY, TR
 AT 329051
 AT 2001-201027
 T 20060615
 JP 2000-80998
 A 20000322
 MARPAT 135:254110

AB A rapid and efficient method is provided for staining and detecting bacteria even in the presence of impurities in a sample (e.g., urine, blood) without culturing it. In this method, a cationic surfactant is added to the sample containing bacteria to promote its dye-permeability. Then, the bacteria is stained with a dye (e.g., fluorescent dye), and detected by flow cytometry.

IC ICM C12Q001-06
ICS C12Q001-04, G01N033-48; C12Q001-06; C12R001-01

CC 9-16 (Biochemical Methods)

ST Section cross-reference(s): 10

IT Bacteria staining fluorescent dye cationic surfactant

IT Surfactants
(cationic; method for staining and detecting bacteria)

IT Cytometry
(flow; method for staining and detecting bacteria)

IT Bacteria (Eubacteria)

IT Blood analysis

IT Buffers

IT Citrobacter freundii

IT Dyes

IT Enterococcus faecalis

IT Escherichia coli

IT Fluorescent dyes

IT Fluorimetry

IT Impurities

IT Klebsiella pneumoniae

IT Light scattering

IT Permeability

IT Pseudomonas

IT Staining, biological

IT Staphylococcus

IT Streptococcus aureus

IT Urine analysis

IT pH

(method for staining and detecting bacteria)

IT Nitrates, biological studies

IT Quaternary ammonium compounds, biological studies

IT Sulfates, biological studies

IT RL: RUU (Biological use, unclassified); BIOL (Biological study); USES

(Uses)

(method for staining and detecting bacteria)

IT 50-21-5, Lactic acid, biological studies 56-40-6, Glycine, biological

studies 60-32-2, 8-Aminocaproic acid 77-92-9D, Citric acid,

salt 88-99-3D, Phthalic acid, salt 107-95-9, β -Alanine

110-15-6, Succinic acid, biological studies 110-17-8, Fumaric acid,

biological studies 6899-10-1 10182-91-9 10182-92-0,

Tetradecyltrimethylammonium 14265-44-2, Phosphate, biological studies

15053-09-5, Decyltrimethylammonium 15461-40-2,

Octadecyltrimethylammonium 24147-36-2 76433-27-7

76433-29-9 150749-57 8 157199-63-8

166196-17-4 361437 94-7 361544-71-0

361544-72-1

RL: RUU (Biological use, unclassified); BIOL (Biological

study); USES (Uses)

(method for staining and detecting bacteria)

IT 24147-36-2 76433-27-7 76433-29-9

150749-57-8 157199-63-8 166196-17-4

361437-94-7 361544-71-0 361544-72-1


RL: RUU (Biological use, unclassified); BIOL (Biological

study); USES (Uses)

(method for staining and detecting bacteria)

RN 24147-36-2 HCAPLUS

CN Quinolinium, 1-methyl-4-[(3-methyl-2-(3H)-benzothiazolylidene)methyl]-, iodide (1:1) (CA INDEX NAME)




RN 76433-27-7 HCAPLUS

CN 3H-Indolium, 2-[4-[4-(dimethylamino)phenyl]-1,3-butadienyl]-1,3,3-trimethyl-, perchlorate (9CI) (CA INDEX NAME)

CM 1

CRN 76433-26-6

CMF C23 H27 N2



CM 2

CRN 14797-73-0

CMF Cl O4



RN 76433-29-9 HCAPLUS

CN Benzothiazolium, 2-[4-[4-(dimethylamino)phenyl]-1,3-butadienyl]-1,3-ethyl-, perchlorate (9CI) (CA INDEX NAME)

10/803,667

August 23, 2007

10/803,667

August 23, 2007

, perchlorate (1:1) (CA INDEX NAME)

CM 1

CRN 76433-28-8

CMF C21 H23 N2 S



CM 2


CRN 14797-73-0

CMF Cl O4



RN 150749-57-8 HCAPLUS


CN Benzothiazolium, 3-[3-(trimethylammonio)propyl]-2-[5-[3-(3-(trimethylammonio)propyl)-2(3H)-benzothiazolylidene]-1,3-pentadienyl]-, tribromide (9CI) (CA INDEX NAME)



● 3 Br-

RN 157199-63-8 HCAPLUS

CN Quinolinium, 4-[3-(3-methyl-2(3H)-benzothiazolylidene)-1-propen-1-yl]-1-[3-(trimethylammonio)propyl]-, iodide (1:2) (CA INDEX NAME)




● 2 I-


RN 166196-17-4 HCAPLUS

CN Quinolinium, 1,1'-[1,3-propanediylbis[(dimethyliminio)-3,1-propanediyl]]bis[4-[3-(3-methyl-2(3H)-benzothiazolylidene)-1-propen-1-yl]]-, iodide (1:4) (CA INDEX NAME)

PAGE 1-A



PAGE 2-A




RN 361437-94-7 HCAPLUS
 CN Benzenesulfonic acid, 4-[[4-[5-(1,3-dibutyltetrahydro-4,6-dioxo-2-thioxo-5(2H)-pyrimidinylidene)-1,3-pentadienyl]-4,5-dihydro-3-methyl-5-oxo-1H-pyrazol-1-yl]]-, compd. with N,N-diethylethanamine (1:2) (9CI) (CA INDEX NAME)

CM 1

CRN 361437-93-6
 CMF C27 H32 N4 O6 S2

PAGE 2-A




CM 2

CRN 121-44-8
 CMF C6 H15 N




RN 361544-71-0 HCAPLUS
 CN Borate(1-), difluoro[2,3,5,6-tetrafluoro-4-sulphophenyl 6-[[[4-(2-[5-[[5-(2-thienyl)-2H-pyrrol-2-ylidene-KN]methyl]-1H-pyrazol-2-yl-KN]ethenyl]phenoxy]acetyl]aminohexanoato(2-)]-], sodium, (T-4) - (9CI) (CA INDEX NAME)


PAGE 1-A



PAGE 1-A




PAGE 1-B



RN 361544-72-1 HCAPLUS
 CN Borate(1-), difluoro[2,3,5,6-tetrafluoro-4-sulphophenyl


6-[[[4-[2-[2-[(2,2'-bi-1H-pyrrol]-5-yl-xyl)methylene]-2H-pyrrol-5-yl-xyl]ethenyl]phenoxy]acetyl]amino]hexanoato(2-)]-, sodium, (T-4)-(9CI) (CA INDEX NAME)

PAGE 1-A



Na⁺

PAGE 1-B




L80 ANSWER 25 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2001:569726 HCAPLUS Full-text
 DOCUMENT NUMBER: 135:149591
 TITLE: Methods and compositions for rapid staining of nucleic acids in whole cells
 INVENTOR(S): Dekka, Chiranjit; Gordon, Kristie M.; Gupta, Ravinder; Horton, Allan
 PATENT ASSIGNEE(S): Coulter International Corp., USA
 SOURCE: U.S., 10 pp.
 CODEN: USXXAM
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|---|------|----------|-----------------|----------|
| US 6271035 | B1 | 20010807 | US 1998-175495 | 19981020 |
| PRIORITY APPLN. INFO.: US 1998-175495 19981020 | | | | |
| AB A rapid fluorescence staining method for facilitating flow cytometry anal. of reticulocytes is described. The method comprises contacting cells with a cocktail containing a detergent, sphering agent, and a cell impermeable dye, such as TO-PRO-3, for about one minute. Advantageously, the inventors have found that the cocktail permits the dye to penetrate the cell membrane rapidly. | | | | |

IC ICM GOIN031-00


INCL 436010000
 CC 9-4 (Biochemical Methods)
 ST compn staining nucleic acid cell
 IT Dyes (cell membrane impermeant; methods and compns. for rapid staining of nucleic acids in whole cells)
 IT Cytometry (flow; methods and compns. for rapid staining of nucleic acids in whole cells)
 IT Staining; biological (fluorescent; methods and compns. for rapid staining of nucleic acids in whole cells)
 IT Biological transport
 Cell
 Cell membrane
 Composition
 Detergents
 Fluorescent dyes
 Mixtures
 Reticulocyte
 Samples
 Solutions
 Spectroscopy
 Spheres
 Stains, biological (methods and compns. for rapid staining of nucleic acids in whole cells)
 DNA
 Nucleic acids
 RNA
 RL: ANT (Analyte); BSU (Biological study, unclassified); ANST (Analytical study); BIOL (Biological study) (methods and compns. for rapid staining of nucleic acids in whole cells)
 IT Detergents (nontonic; methods and compns. for rapid staining of nucleic acids in whole cells)
 IT Laser radiation (red; methods and compns. for rapid staining of nucleic acids in whole cells)
 IT Detergents (zwitterionic; methods and compns. for rapid staining of nucleic acids in whole cells)
 IT 92-32-0, Pyronin Y. 1239-45-8, Ethidium bromide 25535-16-4, Propidium iodide 143413-84-7, TOYO-1 143413-85-8, YOYO-1 152068-09-2, YO-PRO-1 154757-99-0, POPO-3 157199-63-8, TO-PRO-3 161016-55-3, PO-PRO-3 166196-17-4, TOYO-3
 RL: AKG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses) (methods and compns. for rapid staining of nucleic acids in whole cells)
 IT 67-68-5, DMSO, biological studies 9002-93-1, Triton X-100 9036-19-5, Nonidet P-40 55965-84-9, Proclin 300 69227-93-6, Dodecyl-β-D-maltoside
 RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses) (methods and compns. for rapid staining of nucleic acids in whole cells)

IT 157199-63-8, TO-PRO-3 166136 17-4, TOTO-3
 RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses) (methods and compns. for rapid staining of nucleic acids in whole cells)
 RN 157199-63-8 HCAPLUS
 CN Quinolinium, 4-{3-(3-methyl-2(3H)-benzothiazolylidene)-1-propen-1-yl}-[3-(trimethylammonio)propyl]-, iodide (1:2) (CA INDEX NAME)




●2 I-

RN 166196-17-4 HCAPLUS
 CN Quinolinium, 1,1'-[1,3-propanediylbis((dimethyliminio)-3,1-propanediyl)]bis[4-{3-(3-methyl-2(3H)-benzothiazolylidene)-1-propen-1-yl}-, iodide (1:4) (CA INDEX NAME)



PAGE 1-A



REFERENCE COUNT: 25 THERE ARE 25 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L80 ANSWER 26 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2001:284081 HCAPLUS Full-text
 DOCUMENT NUMBER: 134:307569
 TITLE: Microfluidic devices and use of Nernstein voltage sensitive dyes in measuring transmembrane voltage
 INVENTOR(S): Farinas, Javier Anibal; Wada, H. Garrett
 PATENT ASSIGNEE(S): Caliper Technologies Corp., USA
 SOURCE: PCT Int. Appl., 70 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|---|------|----------|-----------------|----------|
| WO 2001027253 | A1 | 20010419 | WO 2000-US27659 | 20001006 |
| W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MN, MW, MX, MY, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW | | | | |
| RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BE, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG | | | | |
| CA 2385618 | A1 | 20010419 | CA 2000-2385618 | 20001006 |
| EP 1222257 | A1 | 20020717 | EP 2000-975224 | 20001006 |
| R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LV, FI, RO, MK, CY, AL | | | | |
| JP 2003511682 | T | 20030325 | JP 2001-530458 | 20001006 |
| US 6537771 | B1 | 20030325 | US 2000-684313 | 20001006 |
| AU 783191 | B2 | 20051006 | AU 2001-13304 | 20001006 |
| US 2004009545 | A1 | 20040115 | US 2003-349396 | 20030121 |
| US 6759191 | B2 | 20040706 | | |

US 2004048239 A1 20040311 US 2003-655697 20030905
 US 6979553 B2 20051227 US 1999-158323P P 19991008
 US 1999-168792P P 19991202
 US 2000-229951P P 20000901
 US 2000-684113 A3 20001006
 WO 2000-US27659 W 20001006
 US 2003-349396 A1 20030121

AB Transmembrane potential measurement methods using cationic dyes, and anionic dyes are provided. Comps. of the cationic and anionic dyes and microfluidic systems which include the dyes and membranes are provided in conjunction with processing elements for transmembrane potential measurements. The time course of SYTO 62 (a cyclic-substituted unsym. cyanine dye) uptake by THP-1 cells depended on transmembrane potential. The changes in the cell transmembrane potential were detected in a microfluidic processor.

IC ICM C12N013-00

CC ICS C12Q001-02; G01N001-30; G01N015-06

IT 9-1 (Biochemical Methods)

Nucleic acids

RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(cationic dye staining; microfluidic devices and use of

Nernstein voltage sensitive dyes in measuring transmembrane voltage)

IT Animal cell

Animal tissue culture

Bacteria (Eubacteria)

Blood cell

Buffers

Cell differentiation

Cell membrane

Chloroplast

Containers

Electric potential

Flow

Fluorometry

Fungi

HeLa cell

Membrane, biological

Membrane potential

Membranes, nonbiological

Microtiter plates

Mitochondria

Plant cell

Plant tissue culture

Sensors

T cell (lymphocyte)

(microfluidic devices and use of Nernstein voltage sensitive dyes in measuring transmembrane voltage)

IT 335080-22-3, RGA 30

RL: ARG (Analytical reagent use); ANST (Analytical

study); USES (Uses)

(RGA 30; microfluidic devices and use of Nernstein voltage sensitive dyes in measuring transmembrane voltage)

IT 335080-22-3, RGA 30

RL: ARG (Analytical reagent use); ANST (Analytical

study); USES (Uses)

(RGA 30; microfluidic devices and use of Nernstein voltage sensitive dyes in measuring transmembrane voltage)

RN 335080-22-3 HCAPJUS

CN Benzenesulfonic acid, 4-[(4-[5-(1,3-dibutylhexahydro-4,6-dioxo-2-thioxo-5-


pyrimidinyl)-2,4-pentadienyldene]-4,5-dihydro-3-methyl-5-oxo-1H-pyrazol-1-yl]-, compd. with N,N-diethylethanamine (1:2) (9CI) (CA INDEX NAME)

CM 1


CRN 118702-42-4

CMF C27 H32 N4 O6 S2

PAGE 1-A



PAGE 2-A



CM 2

CRN 121-44-8

CMF C6 H15 N



REFERENCE COUNT: 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L80 ANSWER 27 OF 46 HCAPLUS COPYRIGHT 2007 ACS ON STN
 ACCESSION NUMBER: 2001:230342 HCAPLUS Full-text
 DOCUMENT NUMBER: 135:207662

TITLE: Multiparameter flow cytometry of bacteria:
 Implications for diagnostics and therapeutics

AUTHOR(S): Shapiro, Howard M.
 CORPORATE SOURCE: West Newton, MA, 02465-2513, USA
 SOURCE: Cytometry (2001), 43(3), 223-226
 CODEN: CYTOOQ; ISSN: 0196-4763

PUBLISHER: Wiley-Liss, Inc.
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB Flow cytometric studies of antibiotic susceptibilities of bacteria have typically measured a single fluorescence parameter, such as membrane potential (indicating viability), or permeability to nucleic acid stains such as propidium (indicating nonviability). Cytometry of bacteria stained simultaneously with a membrane potential dye and a permeability indicator reveals unanticipated complexity. Aliquots of cultures of three bacterial species were stained with the potential-sensitive dye hexamethylindocarbocyanine [DiIc1(3)] and the permeability indicator TO-PRO-3, in the presence and absence of a proton ionophore which collapses the potential gradient. They were analyzed using a dual-laser flow cytometer. Cultures grown under suboptimal conditions appear to contain cells that take up TO-PRO-3 while maintaining membrane potential, although some events showing both high DiIc1(3) fluorescence and high TO-PRO-3 fluorescence may represent clumps. Variations in metabolic patterns between species and within organisms under suboptimal culture conditions or following antibiotic exposure may make it difficult to develop flow cytometric clin. assays for antibiotic susceptibility. However, transient permeabilization of otherwise resistant organisms by sublethal doses of antibiotics may make it possible to treat infections by such organisms with suitably derivatized, otherwise toxic mols.; multiparameter cytometry should be useful in pursuing this approach to therapy.

CC 9-5 (Biochemical Methods)

ST Section cross-reference(s): 1, 10

IT Multiparameter flow cytometry bacteria diagnostic therapeutic

IT Membrane potential
 (biol.; multiparameter flow cytometry of bacteria and implications for diagnostics and therapeutics)

IT Cytometry
 (flow; multiparameter flow cytometry of bacteria and implications for diagnostics and therapeutics)

IT Staining, biological
 (fluorescent; multiparameter flow cytometry of bacteria and implications for diagnostics and therapeutics)

IT Antibiotics
 Bacteria (Eubacteria)

Diagnosis

Fluorescence

Fluorometry

Membrane, biological

Therapy

(multiparameter flow cytometry of bacteria and implications for diagnostics and therapeutics)

Biological transport

IT Implications for diagnostics and therapeutics

(permeation; multiparameter flow cytometry of bacteria and implications for diagnostics and therapeutics)

IT 555-60-2, cccp 25470-94-4 157199-63-8, TO-PRO-3

RU: BUU (biological use, unclassified); BIOL (biological

study); USES (Uses)

(multiparameter flow cytometry of bacteria and implications for diagnostics and therapeutics)


IT 157199-63-8, TO-PRO-3

RL: BUU (biological use, unclassified); BIOL (biological study); USES (Uses)

(multiparameter flow cytometry of bacteria and implications for diagnostics and therapeutics)

RN 157199-63-8 HCAPLUS

CN Quinolium, 4-[3-(3-methyl-2(3H)-benzothiazolylidene)-1-propen-1-yl]-1-[3-(trimethylammonio)propyl]-, iodide (1:2) (CA INDEX NAME)



2 I-

REFERENCE COUNT: 16

16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L80 ANSWER 28 OF 46 HCAPLUS COPYRIGHT 2007 ACS ON STN
 ACCESSION NUMBER: 2001:168247 HCAPLUS Full-text
 DOCUMENT NUMBER: 134:190341
 TITLE: Method and device for counting cells in urine
 INVENTOR(S): Gjelsnes, Oddbjorn; Romning, Oystein
 PATENT ASSIGNEE(S): Optoflow AS, Norway
 SOURCE: PCT Int. Appl., 13 pp.
 CODEN: PIXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|---------------|--|----------|-----------------|----------|
| WO 2001016595 | A1 | 20010308 | WO 2000-NO286 | 20000901 |
| W: | AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KP, KZ, LC, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MX, MY, NZ, OL, OM, OS, PA, PE, PG, PH, PK, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, MD, RU, TJ, TM | | | |
| RW: | GH, GM, KE, LS, MW, NZ, SD, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, | | | |

CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG
 EP 1181553 A1 20020227 EP 2000-959042 20000901
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, MC, IE, SL,
 LT, LV, FI, RO

PRIORITY APPLN. INFO.:

NO 1999-4228 A 19990901
 WO 2000-NO286 W 20000901

AB The invention regards a method and a device for measuring the number of cells in urine. A fixative, a buffer and a dye are added to the urine sample, which is then analyzed in a device for measuring fluorescence.

IC ICM GOIN033-493

CC 9-1 (Biochemical Methods)

ST device counting cell urine

IT Pumps

(Adjustable multichannel; method and device for counting cells in urine)

IT Cytometry

IT (apparatus, flow; method and device for counting cells in urine)

IT Measuring apparatus

(cytometers, flow; method and device for counting cells in urine)

IT Apparatus

Bacteria (Eubacteria)

Buffers

Carriers

Cell

Cell membrane

Cyanine dyes

Dyes

Fluorescent substances

Fluorometers

Fluorometry

Light scattering

Liquids

Mixers (processing apparatus)

Mixing

Pipes and Tubes

Spectrometers

Staining, biological

UV and visible spectroscopy

Urine analysis

(method and device for counting cells in urine)

IT Nucleic acids

RL: BSU (Biological study, unclassified); BIOL (Biological study)

(method and device for counting cells in urine)

IT Biological transport

(permeation; method and device for counting cells in urine)

IT 60-00-4, EDTA, biological studies 64-17-5, Ethanol, biological studies

67-63-0, Isopropanol, biological studies 67-64-1, Acetone, biological

studies 77-86-1, Tris buffer 11129-12-7, Borate 157199-63-8,

TOPRO-3

RL: RUU (Biological use, unclassified); BIOL (Biological

study); USES (Uses)

(method and device for counting cells in urine)

IT 157199-63-8, TOPRO-3

(method and device for counting cells in urine)

RL: RUU (Biological use, unclassified); BIOL (Biological


study); USES (Uses)

(method and device for counting cells in urine)

RN 157199-63-8 HCAPLUS

CN Quinolinium, 4-[3-(3-methyl-2(3H)-benzothiazolylidene)-1-propen-1-yl]-1-[3-

(trimethylammonio)propyl]-, iodide (1:2) (CA INDEX NAME)



● 2 I-

REFERENCE COUNT:

4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L80 ANSWER 29 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2000:688089 HCAPLUS Full-text

DOCUMENT NUMBER: 133:247259

TITLE: Combination of antibiotic and nucleic acid-binding

compound for killing bacteria, including

antibiotic-resistant bacteria

Shapiro, Howard M.

USA

PCT Int. Appl., 31 pp.

CODEN: PIXXD2

Patent

English

1

FAMILY ACC. NUM. COUNT:

PATENT INFORMATION:

PATENT NO.

WO 2000056333

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, PA, PE, PG, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW

RW: GH, GM, KE, LS, MW, SL, SZ, TZ, UG, ZW, AT, BE, CH, CI, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GN, GW, ML, MR, NE, SN, TD, TG

US 6562785

CA 2367149

EP 1165073

EP 1165073

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, CY

JP 2002539259

AU 761609

B2

20030605

JP 2000-606238

AU 2000-39068

20000321

20000321

20000321

20000321

20000321

20000321

20000321

20000321

20000321

20000321

20000321

20000321

20000321

20000321

20000321

20000321

20000321

20000321

20000321

20000321

AT 327754 T 20060615 AT 2000-918218 20000321
 PRIORITY APPLN. INFO.: US 1999-274699 A 19990323
 WO 2000-057500 W 20000321

AB Methods are provided for killing bacteria, including antibiotic-resistant bacteria, by contacting the bacteria with a membrane permeabilizing compound or combination of compds. and a membrane impermeant toxic agent or combination of agents, resulting in the death of the bacteria without substantial injury to the infected host or patient. The invention is also provides related compns. and kits. Further provided are methods of rendering toxic agents, e.g. toxic organic mois., membrane impermeant for use in the methods and compns.

IC ICM A61K031-43
 ICS A61P031-00; A61P033-00

CC 1-5 (Pharmacology)
 Section cross-reference (s): 63
 antibiotic nucleic acid binder bactericide; resistance
 antibiotic nucleic acid binder bactericide

ST Membrane potential
 (biol.; combination of antibiotic and nucleic acid-binding compound for killing bacteria, including antibiotic-resistant bacteria)

IT Antibacterial agents
 Antibiotic resistance
 Antibiotics
 Antimicrobial agents
 Cyanine dyes
 Drug delivery systems
 Drug screening
 Fungicides
 Micrococcus luteus
 Staphylococcus aureus
 (combination of antibiotic and nucleic acid-binding compound for killing bacteria, including antibiotic-resistant bacteria)

IT Nucleic acids
 RL: BSU (Biological study, unclassified); BIOL (Biological study)
 (combination of antibiotic and nucleic acid-binding compound for killing bacteria, including antibiotic-resistant bacteria)

IT Membrane, biological
 (membrane-permeabilizing compds.; combination of antibiotic and nucleic acid-binding compound for killing bacteria, including antibiotic-resistant bacteria)

IT Biological transport
 (permeation; combination of antibiotic and nucleic acid-binding compound for killing bacteria, including antibiotic-resistant bacteria)


IT Cell wall
 (synthesis, inhibitors; combination of antibiotic and nucleic acid-binding compound for killing bacteria, including antibiotic-resistant bacteria)

IT Lactams
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (β-, antibiotics; combination of antibiotic and nucleic acid-binding compound for killing bacteria, including antibiotic-resistant bacteria)

IT Lactams
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(β-, monocyclic; combination of antibiotic and nucleic acid-binding compound for killing bacteria, including antibiotic-resistant bacteria)

IT Antibiotics
 (β-lactam; combination of antibiotic and nucleic acid-binding compound for killing bacteria, including antibiotic-resistant bacteria)
 IT 260-94-6D, Acridine, derivs.
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (and thiazines; combination of antibiotic and nucleic acid-binding compound for killing bacteria, including antibiotic-resistant bacteria)
 IT 56-75-7, Chloramphenicol 57-92-1, Streptomycin, biological studies
 60-54-8, Tetracycline 68-41-7, Cycloserine 114-07-8, Erythromycin 1404-90-6, Vancomycin 1405-87-4, Bacitracin 11111-12-9D, Cephalosporin, derivs. 23686-76-2D, Phenanthridinium, compds. 26787-78-0, Amoxicillin 157199-63-8, TO-PRO-3
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (combination of antibiotic and nucleic acid-binding compound for killing bacteria, including antibiotic-resistant bacteria)
 IT 157199-63-8, TO-PRO-3
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (combination of antibiotic and nucleic acid-binding compound for killing bacteria, including antibiotic-resistant bacteria)
 RN 157199-63-8 HCAPLUS
 CN Quinolium, 4-[3-(3-methyl-2(3H)-benzothiazolylidene)-1-propen-1-yl]-1-[3-(trimethylammonio)propyl]-, iodide (1:2) (CA INDEX NAME)



● 2 I-

REFERENCE COUNT: 9 THERE ARE 9 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L80 ANSWER 30 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2000:553732 HCAPLUS Full-text

DOCUMENT NUMBER: 133.145894
 TITLE: Detection of nucleic acid sequences by amplification as RNA using DNA-dependent RNA polymerase
 INVENTOR(S): Biebricher, Christof K.; Luce, Rudiger; Berendes, Frank; Kessler, Maria; Kalkus, Jutta; Gellersen, Katja; Gottschalk, Gerhard
 PATENT ASSIGNEE(S): Max-Planck-Gesellschaft zur Forderung der Wissenschaften e.V., Germany; Clavigen G.m.b.H.
 SOURCE: PCT Int. Appl., 62 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: German
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|---|------|----------|------------------|----------|
| WO 2000046400 | A1 | 20000810 | WO 2000-EP875 | 20000203 |
| W: CA, JP, US RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE | | | | |
| DE 19904285 | A1 | 20000810 | DE 1999-19904285 | 19990203 |
| EP 1147221 | A1 | 20011024 | EP 2000-905016 | 20000203 |
| R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI | | | | |

PRIORITY APPLN. INFO.:

DE 1999-19904285 A 19990203
 WO 2000-EP875 W 20000203
 AB A method for the qual. or quant. detection of a nucleic acid analyte in a sample by amplification as an RNA using a DNA-dependent RNA polymerase and a probe containing a suitable start site is described. The probe contains a sequence specific to the target sequence and a region that the polymerase uses to start RNA formation from. According to said method the analyte is detected by amplification of the RNA replicon using a DNA-dependent RNA polymerase and subsequent detection of the amplification products. The invention also relates to a nucleic acid which codes for an anal. reagent provided for in the invention and to a test kit for carrying out the above method. The kit also uses a capture probe that can be used to immobilize the target sequence and define an end-point for the amplification product.

ICM C12Q001-68
 ICS C12P019-34
 CC 3-1 (Biochemical Genetics)
 Section cross-reference(s): 9
 IT Bacteriophage SP6
 Coliphage T7
 Enterobacteria phage T3
 (sequence amplification using replication elements of; detection of nucleic acid sequences by amplification as RNA using DNA-dependent RNA polymerase)
 IT 65-61-2, Acridine orange 1239-45-8, Ethidium bromide 24147-36-2, Thiazole orange 25535-16-4, Propidium iodide 152068-09-2, YoPro 1 157199-59-2, ToPro 1
 RL: AFU (Analytical role, unclassified); ANST (Analytical study)
 (as reporter dye; detection of nucleic acid sequences by amplification as RNA using DNA-dependent RNA polymerase)
 IT 24147-36-2, Thiazole orange
 RL: ARU (Analytical role, unclassified); ANST (Analytical study)
 (as reporter dye; detection of nucleic acid sequences by amplification as RNA using DNA-dependent RNA polymerase)

RN 24147-36-2 HCAPLUS
 CN Quinolinium, 1-methyl-4-[(3-methyl-2(3H)-benzothiazolylidene)methyl]-, iodide (1:1) (CA INDEX NAME)



REFERENCE COUNT: 7

THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L80 ANSWER 31 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2000:376371 HCAPLUS Full-text
 DOCUMENT NUMBER: 134:3090

TITLE: In vivo biotinylation studies: specificity of labelling of reticulated platelets by thiazole orange and mepacrine

AUTHOR(S): Robinson, Monique; Machin, Samuel; Mackie, Ian; Harrison, Paul

CORPORATE SOURCE: Department of Haematology, University College

Hospital, London, WC1E 6HX, UK

SOURCE: British Journal of Haematology (2000), 108(4), 859-864

CODEN: BJHEAL; ISSN: 0007-1048

PUBLISHER: Blackwell Science Ltd.


DOCUMENT TYPE: Journal

LANGUAGE: English


AB Animal in vivo biotinylation studies have demonstrated that thiazole orange (TO) labels the youngest cells in the circulation. TO has since been widely used for the measurement of reticulated platelets. As recent findings suggest that at high concns. TO also labels platelet dense granules non-specifically, the value of previous work is unclear. Mepacrine also labels platelet dense granules and can detect storage pool defects. In this study, a mouse in vivo biotinylation model was used to determine the specificity of TO and mepacrine labelling on platelets recently released into the circulation. The mean life span of biotin/TO (low), biotin/TO (high) and mepacrine/TO dual-pos. platelets was 1.4 d (SD 0.5), 2.2 d (SD 0.2) and 2.3 d (SD 0.3) resp. (n = 6) compared with a life span for biotin-pos. platelets of 4.9 d (SD 1.6). TO (low), TO (high) and mepacrine labeled 8.0% (SD 3.1), 43.9% (SD 8.3) and 40.0% (SD 9.9) of the total platelet population resp. (results of 30 samples from six mice), which decreased to 6.8% (SD 3.9), 26.6% (SD 6.9) and 25.7% (SD 10.6) after thrombin degranulation. The shorter life span and lack of thrombin sensitivity of TO (low)-pos. platelets, suggests that TO (low) measures reticulated platelets specifically. The comparative life spans and thrombin sensitivity of TO (high) and mepacrine-pos. platelets suggest that TO (high) labels platelet dense granules. These data also suggest that dense granules are lost during platelet ageing.

CC 13-5 (Mammalian Biochemistry)
ST Section cross-reference(s): 6, 9
platelet reticulated circulation biotin thiazole orange mepacrine staining
IT Biotinylation
Circulation
Fluorescence
Staining, Biological
(in vivo biotinylation studies and specificity of labeling of reticulated platelets by thiazole orange and mepacrine)
IT 58-85-5, Biotin 83-89-6, Mepacrine 107091-89-4, Thiazole orange
RL: APC (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)
(in vivo biotinylation studies and specificity of labeling of reticulated platelets by thiazole orange and mepacrine)
IT 107091-89-4, Thiazole orange
RL: ARS (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)
(in vivo biotinylation studies and specificity of labeling of reticulated platelets by thiazole orange and mepacrine)
RN 107091-89-4 HCAPLUS
CN Quinolifium, 1-methyl-4-[(3-methyl-2(3H)-benzothiazolylidene)methyl]-, 4-methylbenzenesulfonate (1:1) (CA INDEX NAME)

CM 1

CM 2
CRN 24144-08-9
CMP C19 H17 N2 S

CM 2

CRN 16722-51-3
CMP C7 H7 O3 S

REFERENCE COUNT: 21 THERE ARE 21 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L80 ANSWER 32 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 2000-367108 HCAPLUS Full-text
DOCUMENT NUMBER: 133:14302

TITLE: Erythroblast diagnostic flow-cytometry method and reagents

INVENTOR(S): Tsuji, Tomohiro; Sakata, Takashi; Ikeuchi, Yoshiro;

Oguni, Shin'ichiro

PATENT ASSIGNEE(S): Sysmex Corporation, Japan

SOURCE: Eur. Pat. Appl., 39 pp.

CODEN: EPXXDW

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|---|------|----------|-----------------|------------|
| EP 1004880 | A2 | 20000531 | EP 1998-310004 | 19981207 |
| EP 1004880 | A3 | 20030205 | | |
| R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO | | | | |
| JP 2000162209 | A | 20000616 | JP 1998-336916 | 19981127 |
| JP 3886271 | B2 | 20070228 | | |
| US 6664110 | B1 | 20031216 | US 1998-207995 | 19981209 |
| | | | JP 1998-336916 | A 19981127 |

PRIORITY APPLN. INFO.: MARPAT 133:14302

OTHER SOURCE(S):

AB Reagents and a method for simple and rapid discrimination and counting of erythroblasts in peripheral blood or circulatory system-related samples accurately with high precision is disclosed. The reagents include a hemolytic agent for dissolving erythrocytes in a body fluid sample and for conditioning leukocytes and erythroblasts in the sample to be suitable for staining, and including at least one fluorescent dye selected to stain leukocytes and erythroblasts differentially. When the selected fluorescent dye is mixed with the sample, a detectable difference in fluorescence intensity at least between leukocytes and erythroblasts arises under laser illumination in flow cytometric anal. The reagents further include surfactant added to the hemolytic agent, selected to enable flow cytometric discrimination of erythroblasts in the body fluid sample by their maturation stages.

IC ICM G01N033-50

ICS G01N033-58; G01N033-52

CC 9-5 (Biochemical Methods)

Section cross-reference(s): 14

IT Alkyl groups

Amino group

Anions

Blood analysis

Body fluid

Bone marrow

Circulation

Diagnosis

Dissolution

Erythroblast

Erythrocyte

Fluorescent dyes

Fluorometry

Hemolysis

Laser radiation
Leukocyte
Staining, biological
Surfactants
Urine analysis

ph (erythroblast diagnostic flow-cytometry method and reagents)

IT 54-21-7, Sodium salicylate 69-72-7, biological studies 88-99-3, Phthalic acid, biological studies 569-64-2, Malachite green 633-03-4, Brilliant green 3028-99-7, NK-376 3625-57-8, Nile blue a 4727-50-8, Cryptocyanine 18359-88-1, NK-382 20517-94-6, NK-1836 20591-23-5, NK-138 31815-06-0, Sucrose monocrate 33231-00-4, Iodine green 62669-60-7, Oxazine 720 63561-41-1, LD 700 67556-77-8, Oxazine 750 75621-03-3, Chaps 76433-27-7, Lds730 82473-24-3, Chapso 85316-98-9, Mega-8 85618-20-8 85618-21-9 86303-23-3, Decxy-bigchap 89872-07-1, NK-2711 105893-63-8, NK 2825 148565-55-3 178742-72-8, NK-1954

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

IT (erythroblast diagnostic flow-cytometry method and reagents)

76433-27-7, Lds730
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)


RN (erythroblast diagnostic flow-cytometry method and reagents)

CN 3H-Indolium, 2-[4-[(dimethylamino)phenyl]-1,3-butadienyl]-1,3,3-trimethyl-, perchlorate (9CI) (CA INDEX NAME)

CM 1

CRN 76433-26-6

CMF C23 H27 N2



CM 2

CRN 14797-73-0

CMF Cl O4



L80 ANSWER 33 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 1999.746280. HCAPLUS Full-text
DOCUMENT NUMBER: 132:262246
TITLE: Fluorescence enhancement of DNA-bound TO-PRO-3 by incorporation of bromodeoxyuridine to monitor cell cycle kinetics

AUTHOR(S):

Michael .
Beisker, Wolfgang; Weller-Mewe, Eva Maria; Nusse,

CORPORATE SOURCE:

Flow Cytometry Group, GSF-National Research Center for Environment and Health, Neuherberg, 85764, Germany

SOURCE:

Cytometry (1999), 37(3), 221-229

CODEN: CYTODQ; ISSN: 0196-4763

PUBLISHER:

Wiley-Liss, Inc.

DOCUMENT TYPE:

Journal

LANGUAGE:

English

AB

Background: The detection of DNA-incorporated bromodeoxyuridine (BrdUrd) in mammalian cells is a well-known and important technique to study cell cycle. The use of TO-PRO-3 for detection of BrdUrd substitution of DNA by dual-laser flow cytometry has been investigated. Methods: Fluorescence enhancement of TO-PRO-3 in BrdUrd-labeled cells is registered in combination with the fluorescence emission of the intercalating dye propidium iodide (PI) as a total DNA stain to give bivariate DNA/BrdUrd histograms. By the low concentration of only 0.3 µM TO-PRO-3, BrdUrd detection is optimized, and undisturbed total DNA content by PI can be detected as well. TO-PRO-3 is excited by a red HeNe laser and PI by an argon ion laser. Results: In order to understand the binding of TO-PRO-3, energy transfer from PI to TO-PRO-3 has been measured as well as the influence of an external DNA binding dye such as Hoechst 33258 with Adenine-Thymine (AT) binding specificity. Cell cycle studies of human SCL-2 keratinocytes and mouse 3T3 cells prove the method to be as generally applicable as the classical BrdUrd/Hoechst quenching technique, but without need for expensive UV laser excitation. No BrdUrd sensitivity could be found for the similar dyes TO-PRO-1 and YO-PRO-3, whereas TO-PRO-5 and YOYO-3 showed only very little sensitivity to BrdUrd labeling as compared with TO-PRO-3. Conclusions: Cell cycle studies of mammalian cells can be done by dual-laser flow cytometry without the need for UV lasers by using the BrdUrd-dependent fluorescence enhancement of TO-PRO-3. Total DNA content can be measured simultaneously using PI.

CC 9-5 (Biochemical Methods)

ST Section cross-reference(s): 3, 6, 13

IT DNA fluorescence bromodeoxyuridine stain TO PRO 3 cell cycle

(fluorescent; fluorescence enhancement of DNA-bound TO-PRO-3 by incorporation of bromodeoxyuridine to monitor cell cycle kinetics)

IT 59-14-3, Bromodeoxyuridine 23491-45-4, Hoechst 33258 25535-16-4,

Propidium iodide 156312-20-8, YOYO-3 157199-59-2, TO-PRO-1

157199-62-7, YO-PRO-3 157199-63-8, TO-PRO-3 177027-61-1,

TO-PRO-5

RL: AEG (Analytical reagent use); BPR (Biological

process); BSU (Biological study, unclassified); PRP

(Properties); ANST (Analytical study); BIOL (Biological

study); PROC (Process); USES (Uses)

(fluorescence enhancement of DNA-bound TO-PRO-3 by incorporation of

bromodeoxyuridine to monitor cell cycle kinetics)

IT 157199-63-8, TO-PRO-3

RL: AEG (Analytical reagent use); BPR (Biological

process); BSU (Biological study, unclassified); PRP


(Properties); ANST (Analytical study); BIOL (Biological

study); PROC (Process); USES (Uses)

(fluorescence enhancement of DNA-bound TO-PRO-3 by incorporation of

bromodeoxyuridine to monitor cell cycle kinetics)

RN 157199-63-8 HCAPLUS
CN Quinolinium, 4-[3-(3-methyl-2(3H)-benzothiazolylidene)-1-propen-1-yl]-1-[3-(trimethylammonio)propyl]-, iodide (1:2) (CA INDEX NAME)




● 2 I⁻

REFERENCE COUNT: 27 THERE ARE 27 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L80 ANSWER 34 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 1999:504844 HCAPLUS Full-text
DOCUMENT NUMBER: 131:282109
TITLE: Bacterial fingerprinting by flow cytometry:
Bacterial species discrimination
AUTHOR(S): Kim, Jongseong; Jett, James H.; Larson, Erica J.; Penttila, Janetta R.; Marrone, Babetta L.; Keller, Richard A.
CORPORATE SOURCE: Chemical Science and Technology Division, Los Alamos National Laboratory, Los Alamos, NM, 87545, USA
SOURCE: Cytometry (1999), 36(4), 324-332
CODEN: CYTODQ; ISSN: 0196-4763
PUBLISHER: Wiley-Liss, Inc.
DOCUMENT TYPE: Journal
LANGUAGE: English

AB Background: A flow cytometric measurement (FCM) technique has been developed to size DNA fragments. Individual fragments of a restriction digest of genomic DNA, stained with an intercalating dye, are passed through an ultrasensitive cytometer. The measured fluorescence intensity from each fragment is proportional to the fragment length. Methods: The isolation of bacterial genomic DNA and digestion by restriction enzymes were performed inside an agarose plug. Rare cutting enzymes were employed to produce a manageable number of DNA fragments. Electroelution was used to move the DNA fragments from the agarose plug into a solution containing polyamines to protect the DNA from shear-induced breakage. The DNA was stained with the bisintercalating dye thiazole orange homodimer and introduced into our ultrasensitive flow cytometer. A histogram of the fluorescence intensities (fingerprint) was constructed. Results: Gram-pos. *Bacillus globigii* and Gram-neg. *Bacteria Escherichia coli* and *Erwinia herbicola* were distinguished by the fingerprint pattern of restriction fragments of their genomic DNA. DNA sizes determined by FCM are in good agreement with pulsed-field gel electrophoresis (PFGE) anal. Flow cytometry requires only picogram quantities of purified DNA

CC 3-1 (Biochemical Genetics)
ST Section cross-reference(s): 10
IT Bacteria species detn fingerprinting flow cytometry
IT DNA fingerprinting
IT *Bacillus subtilis*
IT *Escherichia coli*
IT Gram-negative bacteria
IT Gram-positive bacteria (Firmicutes)
IT Pantoea agglomerans
IT PFGE (restriction fragment length polymorphism)
IT (bacterial fingerprinting by flow cytometry relating bacterial species discrimination)
IT Cytometry
IT (flow: bacterial fingerprinting by flow cytometry relating bacterial species discrimination)
IT 24147-36-2, Thiazole orange
IT RL: ARU (Analytical role, unclassified); BSU (Biological study, unclassified); ANST (Analytical study); BIOL (Biological study)
IT (bacterial fingerprinting by flow cytometry relating bacterial species discrimination)
IT 24147-36-2, Thiazole orange
IT RL: ARU (Analytical role, unclassified); BSU (Biological study, unclassified); ANST (Analytical study); BIOL (Biological study)
IT (bacterial fingerprinting by flow cytometry relating bacterial species discrimination)
RN 24147-36-2 HCAPLUS
CN Quinolinium, 1-methyl-4-[(3-methyl-2(3H)-benzothiazolylidene)methyl]-, iodide (1:1) (CA INDEX NAME)



● 1 I⁻

REFERENCE COUNT: 25 THERE ARE 25 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L80 ANSWER 35 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 1998:602033 HCAPLUS Full-text
DOCUMENT NUMBER: 129:313077

TITLE:

Nucleic acid (DNA, RNA) quantification and RNA/DNA ratio determination in marine sediments: comparison of spectrophotometric, fluorometric, and high-performance liquid chromatography methods and estimation of detrital DNA

AUTHOR(S):

Dell'anno, A.; Fabiano, M.; Duineveld, G. C. A.; Kok, A.; Danovaro, R.

CORPORATE SOURCE:

Faculty of Science, Marine Science, University of Ancona, Ancona, 60131, Italy

SOURCE:

Applied and Environmental Microbiology (1998), 64 (9), 3238-3245

CODEN: AEMIDF; ISSN: 0099-2240

PUBLISHER:

American Society for Microbiology

DOCUMENT TYPE:

Journal

LANGUAGE:

English

AB

In this study, we compared three methods for extraction and quantification of RNA and DNA from marine sediments: (i) a spectrophotometric method using the diphenylamine assay; (ii) a fluorometric method utilizing selective fluorochromes (thiazole orange for total nucleic acids and Hoechst 33258 for DNA); and (iii) a high-pressure liquid chromatog. (HPLC) method which uses a specific column to sep. RNA and DNA and UV absorption of the nucleic acids for quantification. Sediment samples were collected in the oligotrophic Cretan Sea (eastern Mediterranean, from 40 to 1.540 m in depth) and compared to the distribution and composition of the benthic microbial assemblages (i.e., bacteria and microp Protozoa). DNA concns. measured spectrophotometrically and by HPLC were not significantly different, while fluorometric yields were significantly lower. Such differences appear mainly due to fact that the stain-DNA complex is strongly dependent on the DNA composition and structure. RNA concns. determined by the three methods displayed some differences; fluorometric and spectrophotometric methods obtain RNA concentration by difference and therefore may be biased by DNA ests. By contrast, the HPLC method provides independent assessments of RNA and DNA concns. We tentatively estimated the contribution of the detrital DNA to the total DNA pools in two ways. The two calcs. provided quite similar results indicating that the majority of the DNA pool in the deep-sea sediments was detrital. Microbial RNA generally accounted for almost the entire sedimentary RNA pools below 100-m depth. RNA concns. were found to decrease along the Cretan shelf and slope. The RNA/DNA ratio calculated by using fluorometric DNA concns. was significantly correlated with values of sediment community oxygen consumption only below 100-m depth (dominated by the microbial biomass). These data suggest that the RNA/DNA ratio, based on fluorometric ests. of DNA, can be used as an indicator of benthic metabolic activity, but only when metazoan contribution to the microbial DNA is negligible.

CC

9-16 (Biochemical Methods)
Section cross-reference(s): 61

IT

107091-89-4, Thiazole orange
RL: ANT (Analyte); ANST (Analytical study)

(nucleic acid (DNA, RNA) quantification and RNA/DNA ratio determination in marine sediments: comparison of spectrophotometric, fluorometric, and high-performance liquid chromatog. methods and estimation of detrital DNA)

IT

107091-89-1, Thiazole orange
RL: ANT (Analyte); ANST (Analytical study)

(nucleic acid (DNA, RNA) quantification and RNA/DNA ratio determination in marine sediments: comparison of spectrophotometric, fluorometric, and high-performance liquid chromatog. methods and estimation of detrital DNA)

RN


107091-89-4 HCAPLUS

CN

Quinolium, 1-methyl-4-[(3-methyl-2(3H)-benzothiazolylidene)methyl]-, 4-methylbenzenesulfonate (1:1) (CA INDEX NAME)


CM 1

CRN 24144-08-9
CMF C19 H17 N2 S



CM 2

CRN 16722-51-3
CMF C7 H7 O3 S



REFERENCE COUNT:

50 THERE ARE 50 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE REFORMAT

L80 ANSWER 36 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
1998:365015 HCAPLUS Full-text
ACCESSION NUMBER:
129:38386
DOCUMENT NUMBER:
TITLE:
Method for classifying and counting immature leukocytes using cell hemolysis, staining and flow cytometry

INVENTOR(S):
Sakata, Takashi, Mizukami, Toshihiro; Hatanaka, Kayo
PATENT ASSIGNEE(S):
Toa Medical Electronics Co., Ltd., Japan
SOURCE:
Eur. Pat. Appl., 14 pp.
CODEN: EPXXDW

DOCUMENT TYPE:
Patent
LANGUAGE:
English
FAMILY ACC. NUM. COUNT:
1
PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|---|------|----------|-----------------|------------|
| EP 844481 | A1 | 19980527 | EP 1997-120368 | 19971120 |
| R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO | | | | |
| JP 10206423 | A | 19980807 | JP 1997-289619 | 19971022 |
| US 5958776 | A | 19990928 | US 1997-972103 | 19971117 |
| CN 1183559 | A | 19980603 | CN 1997-123137 | 19971119 |
| PRIORITY APPLN. INFO.: | | | JP 1996-309492 | A 19961120 |

OTHER SOURCE(S): MARPAT 129:38386 JP 1997-289619 A 19971022

AB A flow cytometry method is described for classifying and counting immature leukocytes. The method consists of (1) treating a hematol. sample with a hemolytic agent which maintains immature leukocytes in a viable state and damages other leukocytes, (2) staining the damaged leukocytes with a fluorochrome which can stain damaged cells, and (3) measuring at least one kind of scattered light and at least one kind of fluorescence of the blood cells treated in the preceding step to classify and count leukocytes based on the intensities of the scattered light and the fluorescence. The hemolytic agent contains the following components (1) a polyoxyethylene series nonionic surface active agent for fixing the cytoplasm and cell membrane of immature leukocytes, (2) a solubilizer for damaging the cell membrane of blood cells and shrinking the cells, (3) an amino acid for fixing the cytoplasm and cell membrane of immature leukocytes, and (4) a buffer for making the pH of the resulting solution 5.0 to 9.0 and its osmotic pressure 150 to 600 mOsm/kg. This method can measure immature leukocytes highly precisely, and simultaneously perform the classification of normal leukocytes and the counting of leukocytes.

IC ICM G01N033-50

ICS G01N001-30; G01N033-52

CC 9-5 (Biochemical Methods)

IT Section cross-reference(s): 13

IT Cytometry
(flow; method for classifying and counting immature leukocytes using cell hemolysis, staining and flow cytometry)

IT Staining, biological
Stains, biological
(fluorescent; method for classifying and counting immature leukocytes using cell hemolysis, staining and flow cytometry)

IT Leukocyte
(immature; method for classifying and counting immature leukocytes using cell hemolysis, staining and flow cytometry)

IT Blood analysis
Buffers
Fluorometry
Hemolysis
Laser radiation scattering
Solubilizers
(method for classifying and counting immature leukocytes using cell hemolysis, staining and flow cytometry)

IT Amino acids, analysis
RL: ARU (Analytical role, unclassified); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)
(method for classifying and counting immature leukocytes using cell hemolysis, staining and flow cytometry)

IT Polyoxyalkylene, analysis
RL: ARU (Analytical role, unclassified); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)
(nonionic surface active agent; method for classifying and counting immature leukocytes using cell hemolysis, staining and flow cytometry)

IT Surfactants
(nonionic, polyoxyethylene; method for classifying and counting immature leukocytes using cell hemolysis, staining and flow cytometry)

IT 65282-36-2, Ethidium diazide chloride
RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST

(Analytical study); BIOL (Biological study); USES (Uses)
(ethidium diazide chloride; method for classifying and counting immature leukocytes using cell hemolysis, staining and flow cytometry)

IT 61926-22-5, Ethidium homodimer 1
RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)
(ethidium homodimer-1; method for classifying and counting immature leukocytes using cell hemolysis, staining and flow cytometry)

IT 180389-01-9, Ethidium homodimer 2
RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)
(ethidium homodimer-2; method for classifying and counting immature leukocytes using cell hemolysis, staining and flow cytometry)

IT 1239-45-8, Ethidium bromide 25535-16-4, Propidium iodide 58880-05-0, Ethidium monoazide 68942-32-5, Ethidium-acridine heterodimer 143413-84-7, TOTO-1 157199-59-2, TO-PRO-1 157199-63-8, TO-PRO-3 166196-17-4, TOTO-3
RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)
(method for classifying and counting immature leukocytes using cell hemolysis, staining and flow cytometry)

IT 59-51-8, Methionine 137-16-6, N-Lauroylsarcosine sodium salt 7365-45-9, HEPES 9004-98-2, Polyoxyethylene oleyl ether 25322-68-3D, nonionic surface active agent 189148-50-3
RL: ARU (Analytical role, unclassified); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)
(method for classifying and counting immature leukocytes using cell hemolysis, staining and flow cytometry)

IT 143011-72-7, Granulocyte colony-stimulating factor
RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
(method for classifying and counting immature leukocytes using cell hemolysis, staining and flow cytometry)

IT 157199-63-8, TO-PRO-3 166196-17-4, TOTO-3
RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)
(method for classifying and counting immature leukocytes using cell hemolysis, staining and flow cytometry)


RN 157199-63-8 HCAPLUS
CN Quinolinium, 4-[3-(3-methyl-2(3H)-benzothiazolylidene)-1-propen-1-yl]-1-[3-(trimethylammonio)propyl]-, iodide (1:2) (CA INDEX NAME)

(CH₂)₃-N⁺Me₃

RN 166196-17-4 HCAPLUS

CN Quinolinium, 1,1'-[1,3-propanediyl]bis[(dimethyliminio)-3,1-propanediyl]bis[4-{3-(3-methyl-2(3H)-benzothiazolylidene)-1-propen-1-yl}-, iodide (1:4) (CA INDEX NAME)

PAGE 1-A



IT 189148-50-3

RL: ARU (Analytical role, unclassified); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses) (method for classifying and counting immature leukocytes using cell hemolysis, staining and flow cytometry)


RN 189148-50-3 HCAPLUS

CN Quinolinium, 1-(2-hydroxyethyl)-4-{3-(3-methyl-2(3H)-benzothiazolylidene)-1-propenyl}-, tetrafluoroborate(1-) (9CI) (CA INDEX NAME)

CM 1

CRN 189148-49-0

CMF C22 H21 N2 O S



CM 2

CRN 14874-70-5

CMF B F4

CCI CCS



REFERENCE COUNT:

7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L80 ANSWER 37 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
 1996:577653 HCAPLUS Full-text
 125:216358

TITLE: Spectroscopically detectable nucleic acid ligands

INVENTOR(S): Pitner, James B.; Malinowski, Douglas P.; Vonk, Glenn
 P.; Gold, Larry
 Nexstar Pharmaceuticals, Inc., USA; Becton Dickinson
 and Company

SOURCE: PCT Int. Appl., 36 pp.

CODEN: PIXXDZ


DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 129

PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|------------------------|--|----------|-----------------|-------------|
| WO 9622383 | A1 | 19960725 | WO 1995-US9237 | 19950721 |
| W: | AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT | | | |
| RW: | KE, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GN, ML, MR, NE, SN, TD, TG | | | |
| US 5641629 | A | 19970624 | US 1995-376329 | 19950120 |
| AU 9531401 | A | 19960807 | AU 1995-31401 | 19950721 |
| EP 805870 | A1 | 19971112 | EP 1995-927346 | 19950721 |
| R: | AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE | | | |
| JP 10512751 | T | 19981208 | JP 1995-522232 | 19950721 |
| AU 773741 | B2 | 20040603 | AU 2001-18257 | 20010202 |
| AU 773815 | B2 | 20040610 | AU 2001-29834 | 20010323 |
| AU 2004206993 | A1 | 20040930 | AU 2004-206993 | 20040901 |
| AU 2004206993 | B2 | 20070222 | | |
| AU 2004210606 | A1 | 20041007 | AU 2004-210606 | 20040910 |
| AU 2004242462 | A1 | 20050127 | AU 2004-242462 | 20041223 |
| AU 2004242532 | A1 | 20050127 | AU 2004-242532 | 20041224 |
| PRIORITY APPLN. INFO.: | | | | |
| | | | US 1995-376329 | A 19950120 |
| | | | US 1990-536428 | B2 19900611 |
| | | | AU 1991-82061 | A0 19910610 |
| | | | US 1991-714131 | A3 19910610 |
| | | | US 1992-931473 | A2 19920817 |
| | | | US 1993-134028 | B1 19931007 |
| | | | US 1994-234997 | A2 19940428 |
| | | | US 1994-276271 | A2 19940718 |
| | | | US 1995-443957 | A2 19950518 |
| | | | WO 1995-US9237 | W 19950721 |
| | | | AU 1996-58839 | A3 19960530 |



AU 1996-61611 A3 19960604
 AU 2000-62349 A3 20000724
 AU 2000-67487 A3 20000726
 AU 2001-18257 A 20010202
 AU 2001-29834 A3 20010323

AB The present invention relates to methods of using spectroscopically detectable labeled receptor mols. to determine the presence or absence of a target compound in a sample. The spectroscopic technique may be fluorescence polarization or fluorescence anisotropy. In one embodiment, spectroscopically detectable nucleic acid ligands labeled with fluorescein or thiazole orange are used to determine the presence or absence of biol. targets of interest (e.g., thrombin, elastase, growth factors, choriionic gonadotropin, bacteria, viruses, etc.) in biol. samples (e.g., blood).

IC ICM C12P019-34

ICS C12Q001-68

CC 9-5 (Biochemical Methods)

Section cross-reference(s): 2, 3, 80

IT Animal cell

Bacteria

Blood analysis

Virus

(spectroscopically detectable nucleic acid ligands as receptors in biochem. anal.)

IT 2321-07-SDP, nucleic acid conjugates 107091-39-4DP, Thiazole orange, nucleic acid conjugates 145563-68-4DP, fluorescein derivs. 146159-59-3DP, fluorescein derivs. 169669-13-ODP, fluorescein derivs. 181380-40-SDP, fluorescein derivs. 181593-35-1P 181593-36-2DP, fluorescein derivs.

RL: APG (Analytical reagent use); SPN (Synthetic preparation);

ANST (Analytical study); PREP (Preparation); USES (Uses)

(spectroscopically detectable nucleic acid ligands as receptors in biochem. anal.)

IT 107091-39-4DP, Thiazole orange, nucleic acid conjugates

RL: ARG (Analytical reagent use); SPN (Synthetic preparation);

ANST (Analytical study); PREP (Preparation); USES (Uses)

(spectroscopically detectable nucleic acid ligands as receptors in biochem. anal.)

RN 107091-39-4 HCAPLUS


CN Quinolinium, 1-methyl-4-[(3-methyl-2(3H)-benzothiazolylidene)methyl]-, 4-methylbenzenesulfonate (1:1) (CA INDEX NAME)

CM 1

CRN 24144-08-9

CMF C19 H17 N2 S


CM 2

CRN 16722-51-3
CMF C7 H7 O3 S

L80 ANSWER 38 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 1996:506433 HCAPLUS Full-text
 DOCUMENT NUMBER: 125:162751
 TITLE: Fluorescent viability assay using cyclic-substituted unsymmetrical cyanine dyes
 INVENTOR(S): Millard, Paul J.; Roth, Bruce L.; Yue, Stephen T.; Haugland, Richard P.
 PATENT ASSIGNEE(S): Molecular Probes, Inc., USA
 SOURCE: U.S.: 26 pp., Cont. of U. S. 5,436,134.
 CODEN: USXXAM
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 8
 PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|------------------------|------|----------|--|-------------|
| US 5534416 | A | 19960709 | US 1993-148847 | 19931108 |
| US 5436134 | A | 19950725 | US 1993-90890 | 19930712 |
| US 5545535 | A | 19960813 | US 1993-146328 | 19931101 |
| CA 2133765 | A1 | 19941027 | CA 1994-2133765 | 19940413 |
| CA 2133765 | C | 19991109 | | |
| EP 675924 | A1 | 19951011 | EP 1994-914173 | 19940413 |
| EP 675924 | B1 | 20011212 | | |
| | | | AT, BE, CH, DE, ES, FR, GB, IT, LI, NL | |
| AT 210703 | T | 20011215 | AT 1994-914173 | 19940413 |
| ES 2166777 | T3 | 20020501 | ES 1994-914173 | 19940413 |
| JP 07196930 | A | 19950801 | JP 1994-159824 | 19940712 |
| JP 2005272479 | A | 20051006 | JP 2005-167583 | 20050607 |
| JP 2005344121 | A | 20051215 | JP 2005-167584 | 20050607 |
| JP 2006111884 | A | 20060427 | JP 2005-306416 | 20051020 |
| PRIORITY APPLN. INFO.: | | | US 1993-47883 | B2 19930413 |
| | | | US 1993-90890 | A1 19930712 |
| | | | US 1993-146328 | A2 19931101 |
| | | | US 1993-148847 | A 19931108 |
| | | | WO 1994-USA127 | W 19940413 |
| | | | JP 1994-159824 | A3 19940712 |

OTHER SOURCE(S): MARPAT 125:162751
 GI




AB

The invention relates to a method of analyzing the viability of a sample of cells using an aqueous solution comprising two fluorescent dyes. Dye I has the formula I where R2 is C1-6 alkyl; Z- is a biol. compatible counterion; X is O, S, Se, or NR15, where R15 is H or C1-6 alkyl; or CR16R17, where R16 and R17, which may be the same or different, are independently H or C1-6 alkyl, or the carbons of R16 and R17 taken in combination complete a 5- or 6-membered saturated ring; and the benzazolium is optionally further substituted; n = 0, 1, or 2; Y is CR3:CR4; p and m = 0 or 1, such that p + m = 1; R5 is a C1-6 alkyl; C1-6 alkenyl, C1-6 polyalkenyl, C1-6 alkynyl, or C1-6 polyalkynyl group; or R5 is an OMEGA; R3, R4, R6 and R7, which may be the same or different, are independently H; or a C1-6 alkyl, C1-6 alkenyl, C1-6 polyalkenyl, C1-6 alkynyl or C1-6 polyalkynyl group; or halogen; or OR8, SR8, (NR8R9), where R8 and R9, which may be the same or different, are independently H; or alkyl groups having 1-6 carbons; or 1-2 substituted or unsubstituted alicyclic, heterocyclic, aromatic, or heteroarom. rings, containing 1-4 heteroatoms, wherein the heteroatoms are O, N, or S. R8 and R9 taken in combination are (CH2)2l(CH2)2 where l = 0, NR10, CH2 or a single bond where R10 is H or an alkyl group having 1-6 carbons; or OSO2R19 where R19 is C1-6 alkyl, or C1-6 perfluoroalkyl, or aryl; or an OMEGA; or R6 and R7, taken in combination are (CH2)v where v = 3 or 4, or R6 and R7 form a fused aromatic ring that is optionally further substituted; such that at least one of R3, R4, R5, R6 and R7, or a substituent on the aromatic ring formed by R6 and R7, is an OMEGA; where OMEGA is a cyclic substituent that is attached by a single bond. Fluorescent Dye II selectively stains either viable or nonviable cells with a detectable fluorescent response that is different from the fluorescent response of Dye I. The stained cells are illuminated at a suitable absorption wavelength, and the fluorescent response is detected to distinguish viable and nonviable cells based on the fluorescent response.

IC ICM GOIN033-00
 ICS C12Q001-04; C12Q001-68; C07H001-00
 INCL 436034000
 CC 9-5 (Biochemical Methods)
 ST Section cross-reference(s): 28, 41
 cell viability detn fluorescent dye; stain fluorescent nucleic acid bacteria viability; animal cell viability detn fluorescent dye
 IT Animal cell
 Bacteria
 Cell
 Escherichia coli
 Fibroblast
 Lymphocyte
 Staphylococcus aureus
 (fluorescent cell viability assay using cyclic-substituted unsym. cyanine dyes)
 IT Dyes, cyanine
 Staining, biological
 Stains, biological
 (fluorescent, fluorescent cell viability assay using cyclic-substituted unsym. cyanine dyes)


- IT Bacteria
(gram-neg., fluorescent cell viability assay using cyclic-substituted unsym. cyanine dyes)
- IT Bacteria
(gram-pos., fluorescent cell viability assay using cyclic-substituted unsym. cyanine dyes)
157199-63-8, To-pro-3
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
(TO-PRO 3; fluorescent cell viability assay using cyclic-substituted unsym. cyanine dyes)
- IT 166196-17-4, TO-TO 3
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
(TO-TO 3; fluorescent cell viability assay using cyclic-substituted unsym. cyanine dyes)
- IT 596-09-8, Fluorescein diacetate 1239-45-8, Ethidium bromide 3348-03-6
3546-21-2, Ethidium 24147-36-2, Thiazole orange 25535-16-4, Propidium iodide 36015-30-2, Propidium 61926-22-5, Ethidium homodimer 63783-82-4, Ethidium monozide 104821-25-2 105284-17-1 124412-00-6 12770-45-0 139626-15-6, Tetramethylrhodamine ethyl ester 163831-68-3 169454-17-5 180388-99-2 180389-00-8 180389-01-9
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
(fluorescent cell viability assay using cyclic-substituted unsym. cyanine dyes)
- IT 157199-63-8, To-pro-3
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
(TO-PRO 3; fluorescent cell viability assay using cyclic-substituted unsym. cyanine dyes)
- RN 157199-63-8 HCAPLUS
CN Quinolinium, 4-[3-(3-methyl-2(3H)-benzothiazolylidene)-1-propen-1-yl]-1-[3-(trimethylammonio)propyl]-, iodide (1:2) (CA INDEX NAME)

● 2 I⁻


- IT 166196-17-4, TO-TO 3
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
(TO-TO 3; fluorescent cell viability assay using cyclic-substituted unsym. cyanine dyes)

- RN 166196-17-4 HCAPLUS
CN Quinolinium, 1,1'-[1,3-propanediyl]bis[(dimethyliminio)-3,1'-propanediyl]bis[4-[3-(3-methyl-2(3H)-benzothiazolylidene)-1-propen-1-yl]-, iodide (1:4) (CA INDEX NAME)

PAGE 1-A



PAGE 2-A



- IT 24147-36-2, Thiazole orange
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
(fluorescent cell viability assay using cyclic-substituted unsym. cyanine dyes)
- RN 24147-36-2 HCAPLUS
CN Quinolinium, 1-methyl-4-[(3-methyl-2(3H)-benzothiazolylidene)methyl]-, iodide (1:1) (CA INDEX NAME)

RL: ANT (Analyte); THU (Therapeutic use); ANST (Analytical study); BIOL (Biological study); USES (Uses)
(Use of Na citrate as anticoagulant before thiazole orange staining of platelet RNA as marker for platelet production in thrombocytopenic)

IT

Blood platelet
(disease, thrombocytopenia, use of Na citrate as anticoagulant before thiazole orange staining of platelet RNA as marker for platelet production in thrombocytopenic)

IT

68-04-2, Sodium citrate
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(anticoagulant; use of Na citrate as anticoagulant before thiazole orange staining of platelet RNA as marker for platelet production in thrombocytopenic)

IT

107091-89-4, Thiazole orange
RL: ARG (Analytical reagent use); THU (Therapeutic use)
; ANST (Analytical study); BIOL (Biological study);
USES (Uses)

(Use of Na citrate as anticoagulant before thiazole orange staining of platelet RNA as marker for platelet production in thrombocytopenic)

IT

107091-89-4, Thiazole orange
RL: ARG (Analytical reagent use); THU (Therapeutic use)
; ANST (Analytical study); BIOL (Biological study);
USES (Uses)


(Use of Na citrate as anticoagulant before thiazole orange staining of platelet RNA as marker for platelet production in thrombocytopenic)

RN

107091-89-4 HCAPLUS
CN Quinolinium, 1-methyl-4-[(3-methyl-2(3H)-benzothiazolylidene)methyl]-, 4-methylbenzenesulfonate (1:1) (CA INDEX NAME)


CM 1

CRN 24144-08-9
CMF C19 H17 N2 S



CM 2

CRN 16722-51-3
CMF C7 H7 O3 S



● I -

L80 ANSWER 39 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 1996:455535 HCAPLUS Full-text
DOCUMENT NUMBER: 125:216038

TITLE: Improved method of staining RNA in platelets for the evaluation of platelet production in thrombocytopenic patients

AUTHOR(S): Yamabe, Kozue; Satoh, Sachiko; Tsukada, Toshiyasu

CORPORATE SOURCE: Dep. Hematol. Lab., Toranomon Hosp., Tokyo, 105, Japan

SOURCE: Rinsho Byori (1996), 44(7), 681-686

CODEN: RBYOAI; ISSN: 0047-1860

PUBLISHER: Rinsho Byori Kankokai

DOCUMENT TYPE: Journal

LANGUAGE: Japanese

AB Measurement of RNA stained platelets was proven to be an easy and useful laboratory test to evaluate the state of platelet production in the bone marrow. As the normal value of platelet with thiazole orange (TO)-stained RNA in the original method was low, the values of TO-stained platelets in the cases with platelet hypoproduction were within the normal range, but ideally, they should be below the normal range. We modified the original method by using Na citrate as the anticoagulant instead of EDTA, and by keeping the TO-stained preparation at 4° until the fluorescence was measured. The normal value of TO-stained platelets was elevated to $22.5 \pm 3.3\%$ ($n = 40$, $M \pm 1SD$) or $54 \pm 10 + 109/L$. Twenty-seven out of 40 thrombocytopenic cases with ITP (idiopathic thrombocytopenic purpura) showed an elevated percentage of TO-stained platelets, 11 cases showed a normal and only 2 cases showed a percentage lower than normal. By contrast, 9 out of 12 cases with platelet hypoproduction showed a lower percentage of TO-stained platelets and no cases showed a value higher than normal. The sensitivity and specificity of this modified RNA staining method for distinguishing thrombocytopenic cases with platelet hyperdestruction from that with hypoproduction were 96% and 75%, resp.

CC 9-4 (Biochemical Methods)

ST Section cross-reference(s): 14


platelet RNA staining thiazole orange thrombocytopenia; sodium citrate platelet RNA staining thrombocytopenia

IT Blood platelet

Staining, biological

(use of Na citrate as anticoagulant before thiazole orange staining of platelet RNA as marker for platelet production in thrombocytopenic)

IT Ribonucleic acids



L80 ANSWER 40 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 1996-443964 HCAPLUS Full-text
DOCUMENT NUMBER: 125-81256

TITLE: permeability as fluorescent stains for nucleic acids
permeability

INVENTOR(S): Yue, Stephen T.; Singer, Victoria L.; Roth, Bruce L.;
Moser, Thomas J.; Millard, Paul J.; Jones, Laurie J.;
Jin, Xiaokui; Haugland, Richard P.; Poot, Martin
Molecular Probes, Inc., USA
PCT Int. Appl., 85 pp.
CODEN: PIXXD2

PATENT ASSIGNEE(S):
SOURCE:

DOCUMENT TYPE: Patent
LANGUAGE: English

FAMILY ACC. NUM. COUNT: 8
PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|--|------|----------|-----------------|-------------|
| WO 9613552 | A2 | 19960509 | WO 1995-US13706 | 19951027 |
| WO 9613552 | A3 | 19960711 | | |
| W: AU, CA, JP | | | | |
| DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE | | | | |
| US 5658751 | A | 19970819 | US 1994-331031 | 19941027 |
| CA 2179284 | A1 | 19960509 | CA 1995-2179284 | 19951027 |
| CA 2179284 | C | 20060404 | | |
| AU 9539672 | A | 19960523 | AU 1995-39672 | 19951027 |
| AU 714890 | B2 | 20000113 | | |
| EP 740689 | A1 | 19961106 | EP 1995-937613 | 19951027 |
| EP 740689 | B1 | 20020130 | | |
| R: AT, BE, CH, DE, FR, GB, LI, NL | | | | |
| JP 09507879 | T | 19970812 | JP 1996-514689 | 19951027 |
| AT 212653 | T | 20020215 | AT 1995-937613 | 19951027 |
| PRIORITY APPL. INFO.: | | | US 1994-331031 | A 19941027 |
| | | | US 1993-47683 | B2 19930413 |
| | | | US 1994-90890 | A2 19940712 |
| | | | WO 1995-US13706 | W 19951027 |

OTHER SOURCE(S): MARPAT 125-81256
AB The invention describes the preparation and use of fluorescent stains for nucleic acids derived from unsym. cyanine dyes comprising a substituted benzazolum ring system linked by a methine bridge to a pyridinium or quinoxalinium ring system. The cyanine dyes of the invention possess a high sensitivity to oligonucleotides and larger nucleic acid polymers in a wide range of cells and gels, and are useful for the anal. of cell structure, membrane integrity or function, and determination of cell cycle distribution.

IC ICM C09B023-02
CC 9-4 (Biochemical Methods)
ST Section cross-reference(s): 3, 13, 14, 41
ST nucleic acid detection cyanine fluorescent stain; dye cyanine
nucleic acid stain; animal cell organelle stain
cyanine dye; cancer cell stain fluorescent cyanine dye
IT Salmonella typhimurium
(metabolic activity; substituted unsym. cyanine dyes with selected

IT permeability as fluorescent stains for nucleic acids)

Animal cell
Animal tissue
Animal tissue culture
Antibiotics
Bacteria
Bactericides, Disinfectants, and Antiseptics
Blood analysis
Body fluid
Cell
Cell cycle
Cell membrane
Cell nucleus
Cell proliferation
Chromosome
Cytoplasm
Dyes, cyanine
Eukaryote
Food analysis
Genetic polymorphism
Lymphocyte
Macrophage
Microorganism
Microorganism metabolism
Mitochondria
Monocyte
Neutrophil
Nucleoid
Parasite
Pharmaceutical analysis
Polymerase chain reaction
Saccharomycetes cerevisiae
Viroid
Virus
Yeast

(substituted unsym. cyanine dyes with selected permeability as fluorescent stains for nucleic acids)
Deoxyribonucleic acids
Nucleic acids
Proteins, analysis
Ribonucleic acids
Ribonucleic acids, ribosomal
RL: ANT (Analyte); ANST (Analytical study)
(substituted unsym. cyanine dyes with selected permeability as fluorescent stains for nucleic acids)
Agglutinins and Lectins
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
(substituted unsym. cyanine dyes with selected permeability as fluorescent stains for nucleic acids)
Antibodies
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
(substituted unsym. cyanine dyes with selected permeability as fluorescent stains for nucleic acids)
Enzymes
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
(substituted unsym. cyanine dyes with selected permeability as fluorescent stains for nucleic acids)
Pyridinium compounds
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)

(substituted unsym. cyanine dyes with selected permeability as fluorescent stains for nucleic acids)

IT Animal cell line
(3T3, substituted unsym. cyanine dyes with selected permeability as fluorescent stains for nucleic acids)

IT Animal cell line
(A431, substituted unsym. cyanine dyes with selected permeability as fluorescent stains for nucleic acids)

IT Enzymes
RL: ANT (Analyte); ANST (Analytical study)
(DNA-supercoiling, substituted unsym. cyanine dyes with selected permeability as fluorescent stains for nucleic acids)

IT Animal cell line
(MDCK, substituted unsym. cyanine dyes with selected permeability as fluorescent stains for nucleic acids)

IT Animal cell line
(P3X, substituted unsym. cyanine dyes with selected permeability as fluorescent stains for nucleic acids)

IT Meat
(beef, substituted unsym. cyanine dyes with selected permeability as fluorescent stains for nucleic acids)

IT Onium compounds
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
(benzazolum, substituted unsym. cyanine dyes with selected permeability as fluorescent stains for nucleic acids)

IT Cytochrome
(flow, substituted unsym. cyanine dyes with selected permeability as fluorescent stains for nucleic acids)

IT Dyes
Staining, biological
Stains, biological
(fluorescent, substituted unsym. cyanine dyes with selected permeability as fluorescent stains for nucleic acids)

IT Electrophoresis and ionophoresis
(gel, substituted unsym. cyanine dyes with selected permeability as fluorescent stains for nucleic acids)

IT Nucleotides, analysis
RL: ANT (Analyte); ANST (Analytical study)
(oligo-, substituted unsym. cyanine dyes with selected permeability as fluorescent stains for nucleic acids)

IT Biological transport
(permeation, substituted unsym. cyanine dyes with selected permeability as fluorescent stains for nucleic acids)

IT Onium compounds
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
(quinolinium, substituted unsym. cyanine dyes with selected permeability as fluorescent stains for nucleic acids)

IT Organelle
(vacuole, substituted unsym. cyanine dyes with selected permeability as fluorescent stains for nucleic acids)

IT 9001-99-4, RNase 9003-98-9, DNase 9075-08-5 24937-83-5, Poly rA 24938-09-1, Poly dA-poly dT 25086-81-1 25191-14-4 25191-20-2, Poly dA 25512-84-9, Poly dG-poly dC 25609-92-1, Poly dC 27416-86-0, Poly rU 27732-54-3, Poly dI 30811-80-4 37228-74-3, Exonuclease 80449-01-0, Topoisomerase
RL: ANT (Analyte); ANST (Analytical study)
(substituted unsym. cyanine dyes with selected permeability as fluorescent stains for nucleic acids)

IT 951-78-0D, Deoxyuridine, halogenated 23491-45-4, HOE33258
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)

(substituted unsym. cyanine dyes with selected permeability as fluorescent stains for nucleic acids)

IT 1239-45-8, Ethidium bromide 23491-52-3, HOE33342 24144-08-9 143413-84-7, TOTO-1 143413-86-9, Oxazole yellow 157199-59-2, TO-PRO-1 173080-70-1, SYTO 14
RL: ARG (Analytical reagent use); PRP (Properties); ANST (Analytical study); USES (Uses)
(substituted unsym. cyanine dyes with selected permeability as fluorescent stains for nucleic acids)

IT 161057-69-8
RL: ARG (Analytical reagent use); RCT (Reactant); ANST (Analytical study); RACT (Reactant or reagent); USES (Uses)
(substituted unsym. cyanine dyes with selected permeability as fluorescent stains for nucleic acids)

IT 2540-30-9P 161057-94-9P
RL: ARG (Analytical reagent use); RCT (Reactant); SPN (Synthetic preparation); ANST (Analytical study); PREP (Preparation); RACT (Reactant or reagent); USES (Uses)
(substituted unsym. cyanine dyes with selected permeability as fluorescent stains for nucleic acids)


IT 98251-83-9P 178918-68-8P 178918-69-9P 178918-70-2P 178918-71-3P 178918-72-4P 178918-73-5P 178918-74-6P 178918-75-7P 178918-76-8P 178918-77-9P 178918-78-0P 178918-79-1P 178918-80-4P 178918-81-5P 178918-82-6P 178918-83-7P 178918-84-8P 178918-85-9P 178918-86-0P 178918-87-1P 178918-88-2P 178918-89-3P 178918-90-6P 178918-91-7P 178918-92-8P 178918-93-9P 178918-94-0P 178918-95-1P 178918-96-2P 178918-97-3P 178918-98-4P 178918-99-5P 178919-00-1P 178919-01-2P 178919-02-3P 178919-04-5P 178919-05-6P 178919-06-7P 178919-07-8P 178919-08-9P 178919-09-0P 178919-10-3P 178919-11-4P 178919-12-5P 178919-13-6P 178919-14-7P 178919-15-8P 178919-16-9P 178919-17-0P 178919-18-1P 178919-19-2P 178919-20-5P 178919-21-6P 178919-22-7P 178919-23-8P 178919-24-9P 178919-31-8P 220222-55-9P
RL: ARG (Analytical reagent use); SPN (Synthetic preparation); ANST (Analytical study); PREP (Preparation); USES (Uses)
(substituted unsym. cyanine dyes with selected permeability as fluorescent stains for nucleic acids)

IT 69-53-4, Ampicillin
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)
(substituted unsym. cyanine dyes with selected permeability as fluorescent stains for nucleic acids)

IT 74-88-4, Methyl iodide, reactions 108-02-1, 2-Dimethylaminoethanol 108-47-4, 2,4-Lutidine 607-66-9, 2-Hydroxy-4-methylquinoline 627-31-6, 1,3-Diiodopropane 696-62-8, 4-Iodoanisole 927-58-2, 4-Bromobutyl chloride 1193-02-8, 4-Aminothiophenol 1198-37-4, 2,4-Dimethylquinoline 2349-67-9, 5-Amino-1,3,4-thiadiazole-2-thiol 2584-47-6 6760-40-3, 1,2-Dimethyl-4-quinolone 10025-87-3, Phosphorus oxychloride 19475-28-6 22049-05-4, 1,2-Dimethyl-4-methoxyquinolinium iodide 39759-82-5 55514-14-2, 3-Methyl-2-methylthiobenzothiazolium tosylate 57876-69-4, 2-Chloro-3-methylquinoline 6104-90-3 81287-35-6 127527-22-4, 4-(Dimethylamino)butyl chloride 178919-25-0 178919-28-3 178919-32-9
RL: RCT (Reactant); RACT (Reactant or reagent)
(substituted unsym. cyanine dyes with selected permeability as fluorescent stains for nucleic acids)

IT 18820-83-2P, Pyridinium iodide 178919-26-1P 178919-27-2P 178919-29-4P 178919-30-7P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(substituted unsym. cyanine dyes with selected permeability as

IT 14341: 8⁺, Oxazole yellow
 RL: ARG (Analytical reagent use); PRP (Properties); ANST (Analytical study); USES (Uses)
 (substituted unsym. cyanine dyes with selected permeability as fluorescent stains for nucleic acids)
 RN 143413-86-9 HCAPLUS
 CN Quinolinium, 1-methyl-4-[(3-methyl-2(3H)-benzoxazolylidene)methyl]-, iodide (1:1) (CA INDEX NAME)



• I:

L80 ANSWER 41 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STW
 ACCESSION NUMBER: 1996:367649 HCAPLUS Full-text
 DOCUMENT NUMBER: 125:81301
 TITLE: Reagent and method for analyzing solid components in urine
 INVENTOR(S): Inoue, Junya
 PATENT ASSIGNEE(S): Toa Medical Electronics Co., Ltd., Japan
 SOURCE: Eur. Pat. Appl., 30 pp.
 CODEN: EPXDXM
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|-------------|------|--------------------------------|-----------------|----------|
| EP 708334 | A2 | 19960424 | EP 1995-610053 | 19951019 |
| EP 708334 | A3 | 19960918 | | |
| EP 708334 | B1 | 20010523 | | |
| | R: | CH, DE, ES, FR, GB, IT, LI, NL | | |
| JP 08170960 | A | 19960702 | JP 1995-267454 | 19951016 |
| JP 3580615 | B2 | 20041027 | | |
| CA 2160962 | A1 | 19960421 | CA 1995-2160962 | 19951019 |
| AU 9534366 | A | 19960502 | AU 1995-34366 | 19951019 |
| AU 701948 | B2 | 19990211 | | |
| EP 1089078 | A1 | 20010404 | EP 2000-123791 | 19951019 |
| EP 1089078 | B1 | 20070228 | | |
| | R: | CH, DE, ES, FR, GB, IT, LI, NL | | |
| ES 2156927 | T3 | 20010801 | ES 1995-610053 | 19951019 |
| US 5891733 | A | 19990406 | US 1995-545939 | 19951020 |
| | | | JP 1994-255580 | 19941020 |

PRIORITY APPLN. INFO.:

OTHER SOURCE(S): MARPAT 125:81301
 AB A reagent for analyzing solid components in urine comprising: (i) a buffer agent for maintaining pH at 5.0 to 9.0, (ii) an osmotic pressure compensating agent for maintaining osmotic pressure at 100 mOsm/kg to 600 mOsm/kg, (iii) a first dye which is a condensed benzene derivative, (i.v.) a second fluorescent dye capable of staining a damaged cell, and (v) a chelating agent. A diluent solution and a dyeing solution were prepared from pH 7.0 50 mM HEPES, sodium propionate (in an amount to adjust osmotic pressure at 150 mOsm/kg), and EDTA tri-K salt 0.4% and a dyeing solution consisting of 400 ppm 1st dye, and 1600 ppm second fluorescent dye.
 IC ICM G01N033-50
 ICS G01N033-569
 ICA G01N015-14; C12Q001-04
 CC 9-15 (Biochemical Methods)
 ST urine analysis chelating osmotic dye
 IT Chelating agents
 Erythrocyte
 Osmotic pressure
 Urine analysis
 (reagent composition containing dyes for analyzing solid components in urine)
 IT Dyes
 RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
 (fluorescent, reagent composition containing dyes for analyzing solid components in urine)
 IT 514-73-8, NK-136
 RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
 (NK 136; reagent composition containing dyes for analyzing solid components in urine)
 IT 20591-23-5, NK-138
 RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
 (NK 138; reagent composition containing dyes for analyzing solid components in urine)
 IT 15185-43-0, NK-1511
 RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
 (NK 1511; reagent composition containing dyes for analyzing solid components in urine)
 IT 3071-69-0, NK 1590
 RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
 (NK 1590; reagent composition containing dyes for analyzing solid components in urine)
 IT 20517-94-6, NK-1836
 RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
 (NK 1836; reagent composition containing dyes for analyzing solid components in urine)
 IT 178742-72-8
 RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
 (NK 1954; reagent composition containing dyes for analyzing solid components in urine)
 IT 89872-07-1
 RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
 (NK 2711; reagent composition containing dyes for analyzing solid components in urine)

components in
urine)

IT 76433-27-7
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
(NK 2780; reagent composition containing dyes for analyzing solid components in urine)

IT 76433-29-9
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
(NK 2783; reagent composition containing dyes for analyzing solid components in urine)

IT 2642-25-3, NK-321
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
(NK 321; reagent composition containing dyes for analyzing solid components in urine)

IT 66230-26-0
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
(NK 375; reagent composition containing dyes for analyzing solid components in urine)

IT 3028-99-7
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
(NK 376; reagent composition containing dyes for analyzing solid components in urine)

IT 36536-22-8, NK-529
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
(NK 529; reagent composition containing dyes for analyzing solid components in urine)

IT 52181-10-9
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
(NK 96; reagent composition containing dyes for analyzing solid components in urine)

IT 62669-60-7, Oxazine 720
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
(Oxazine 720; reagent composition containing dyes for analyzing solid components in urine)

IT 85256-40-2
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
(Oxazine 750 perchlorate; reagent composition containing dyes for analyzing solid components in urine)

IT 14969-56-3
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
(Rhodanile blue; reagent composition containing dyes for analyzing solid components in urine)

IT 3521-06-0, Basic blue 1
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
(Basic blue 1; reagent composition containing dyes for analyzing solid components in urine)

IT 569-64-2, Basic green 4
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
(Basic green 4; reagent composition containing dyes for analyzing solid components in urine)

IT 633-03-4
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
(Basic green; reagent composition containing dyes for analyzing solid components in urine)

IT 60786-96-1
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
(Oxazine 4; reagent composition containing dyes for analyzing solid components in urine)

IT 82-94-0 1934-16-3, Basic blue 24 2381-85-3, Nile Blue chloride 1799-02-2, Capri Blue GON 17572-97-3, Tripotassium EDTA 33231-00-4, Iodine green 89106-91-2, Basic blue 124 177772-75-7, Capri Blue BB 3H-Indolium, 2-[4-(4-(dimethylamino)phenyl)-1,3-butadienyl]-1,3,3-trimethyl-, perchlorate (9CI) (CA INDEX NAME)


IT 76433-27-7
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
(NK 2780; reagent composition containing dyes for analyzing solid components in urine)

RN 76433-27-7 HCAPLUS
CN 3H-Indolium, 2-[4-(4-(dimethylamino)phenyl)-1,3-butadienyl]-1,3,3-trimethyl-, perchlorate (9CI) (CA INDEX NAME)

CM 1
CRN 76433-26-6
CMF C23 H27 N2

CM 2
CRN 14797-73-0
CMF C1 O4

IT 76433-29-9
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)



study); USES (Uses)
(NK 2783; reagent composition containing dyes for analyzing solid components in urine)

RN 76433-29-9 HCAPLUS
CN Benzothiazolium, 2-[4-(4-(dimethylamino)phenyl)-1,3-butadien-1-yl]-3-ethyl-1-perchlorate (1:1) (CA INDEX NAME)

CM 1

CRN 76433-28-8

CMF C21 H23 N2 S



CM 2


CRN 14797-73-0

CMF Cl O4



IT 2642-25-3, NK-321
RL: ARG (analytical reagent use); ANST (Analytical study); USES (Uses)
(NK 321; reagent composition containing dyes for analyzing solid components in urine)

RN 2642-25-3 HCAPLUS
CN Quinolinium, 1-ethyl-4-[3-(3-ethyl-2(3H)-benzothiazolylidene)-1-propenyl]-, iodide (9CI) (CA INDEX NAME)



• I •

L80 ANSWER 42 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 1996:83229 HCAPLUS Full-text
DOCUMENT NUMBER: 124:280288

TITLE: Comparison of self-sustained sequence-replication reaction systems
AUTHOR(S): Gebinoga, Michael; Oehlenschlaeger, Frank
CORPORATE SOURCE: Inst. for Molecular Biotechnology, Jena, Germany
SOURCE: European Journal of Biochemistry (1996), 235(1/2), 256-61
CODEN: EJBICAI; ISSN: 0014-2956

PUBLISHER: Springer
DOCUMENT TYPE: Journal
LANGUAGE: English

AB The 3SR (self-sustained sequence-replication) reaction is a very efficient method for isothermal amplification of target DNA or RNA sequences in vitro. This method requires 3 enzymic activities: reverse transcriptase, DNA-dependent RNA polymerase and Escherichia coli RNase H. The original protocol was modified by using human immunodeficiency virus (HIV)-1 reverse transcriptase instead of avian myeloblastosis virus (AMV) reverse transcriptase to allow amplification with T7 RNA polymerase but without E. coli RNase H. Comparison of the incorporation kinetics between the conventional 3-enzyme 3SR and the 2-enzyme 3SR shows differences in the kinetic behavior. Furthermore, by the new 2-enzyme 3SR, the amplified RNA is obtained in a purer form compared with the expts. with 3-enzyme 3SR. 3SR should be adapted as a useful tool for Darwinian evolutionary expts.

CC 3-1 (Biochemical Genetics)

ST self sustained sequence replication reverse transcriptase;
nucleic acid amplification fluorescence detection

IT Genetic methods

(3SR (self-sustained sequence-replication); 2 enzyme 3SR
system using human immunodeficiency virus-1 reverse transcriptase and
phage T7 RNA polymerase compared to 3 enzyme 3SR)

IT Virus, bacterial


(T7, RNA polymerase; 2 enzyme 3SR system using human immunodeficiency
virus-1 reverse transcriptase and phage T7 RNA polymerase compared to 3
enzyme 3SR)

IT 143413-84-7

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES
(Uses)

(TOTO; as label for self-sustained sequence-replication
reaction systems)

IT 24147-36-2, Thiazole orange
 RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)
 (Thiazole orange; as label for self-sustained sequence-replication reaction systems)
 IT 24147-36-2, Thiazole orange
 RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)
 (Thiazole orange; as label for self-sustained sequence-replication reaction systems)
 RN 24147-36-2 HCAPLUS
 CN Quinolium, 1-methyl-4-[(3-methyl-2(3H)-benzothiazolylidene)methyl]-, iodide (1:1) (CA INDEX NAME)



L80 ANSWER 43 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 1995:623505 HCAPLUS Full-text
 DOCUMENT NUMBER: 124:4485
 TITLE: Fluorescence energy transfer and intramolecular energy transfer in particles using novel compounds
 INVENTOR(S): Buechler, Kenneth Francis; Noar, Joseph Barry; Tadesse, Lema
 PATENT ASSIGNEE(S): Biosite Diagnostics Inc., USA
 SOURCE: PCT Int. Appl., 138 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 7
 PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|---|------|----------|-----------------|----------|
| WO 9508772 | A1 | 19950330 | WO 1994-US10826 | 19940923 |
| W: AU, CA, JP | | | | |
| RM: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE | | | | |
| US 6238931 | B1 | 20010529 | US 1994-274534 | 19940712 |
| CA 2149419 | A1 | 19950330 | CA 1994-2149419 | 19940923 |
| CA 2149419 | C | 20070515 | | |
| AU 9480112 | A | 19950410 | AU 1994-80112 | 19940923 |
| EP 670041 | A1 | 19950906 | EP 1994-931287 | 19940923 |
| EP 670041 | B1 | 20020130 | | |
| R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE | | | | |


JP 08503994 T 19960430 JP 1995-509970 19940923
 JP 3773949 B2 20060510
 AT 212721 T 20020215 AT 1994-931287 19950330
 US 2002061602 A1 20020523 US 2001-776599 20010201
 US 7083984 B2 20060801
 PRIORITY APPLN. INFO.:
 US 1993-126367 A 19930924
 US 1993-118708 A 19931018
 US 1994-274534 A 19940712
 US 1994-311098 A2 19940923
 WO 1994-US10826 W 19940923
 US 1995-409298 A2 19950323
 US 1995-409825 B2 19950323
 US 1996-620597 A1 19960322
 US 1998-66255 A2 19980424
 AB Particles and methods are disclosed for the detection or visualization of analytes, including nucleic acids by using fluorescence energy transfer or intramol. energy transfer. Particles comprising an energy donor as a first component and a fluorescent dye as a second component positioned in said particles at an energy exchanging distance from one another, wherein the two components have a Stokes shift of 250 nm, said particle having bound on its surface, a protein, polypeptide, nucleic acid, nucleotide or protein containing ligand analog are disclosed and claimed. In addition, novel fluorescent dyes are described which exhibit intramol. energy transfer for use in labeling various mols., proteins, polypeptides, nucleotides and nucleic acids or incorporating into particles. Many novel phthalocyanine derivs. and hybrid phthalocyanine derivs. are disclosed and claimed. Such derivs. also may contain an electron transfer subunit. Axial ligands may be covalently bound to the metals contained in the hybrid phthalocyanine derivs. Numerous compds. capable of intramol. energy transfer as well as compds. for fluorescence energy transfer are claimed.
 IC ICM G01N033-546
 CCS C09B047-04
 CC 9-5 (Biochemical Methods)
 IT Section cross-reference(s): 15, 41, 74, 80
 Blood analysis
 Colloids
 Fluorescence quenching
 Fluorescent substances
 Fluorometers
 Immunocassay
 Latex
 Liposome
 Urine analysis
 (fluorescence and intramol. energy transfer in particles for biochem. anal.)
 IT 514-73-8 519-62-0, Chlorophyll b 2321-07-5 3071-70-3 14806-50-9
 16595-48-5 17094-16-5 23481-50-7 24796-94-9, Oxazine 1 perchlorate
 30753-88-9 53213-94-8 53655-17-7 56089-72-6 70365-30-9
 83484-76-8 86880-07-1 94052-41-2 97148-81-7 97807-64-2
 116453-73-7 122711-10-8 150749-57-8 163968-80-7
 163968-81-8 163968-82-9 163968-84-1 163968-85-2 163968-86-3
 163968-87-4 163968-88-5 163968-89-6 163968-90-9 163968-91-0
 163968-92-1 163968-93-2 163968-95-4 163969-13-9 163969-14-0
 164106-16-5 17118-91-5 17118-92-6D, reaction with silicon phthalocyanine 17118-93-7 17118-99-3
 RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)
 (fluorescence and intramol. energy transfer in particles for biochem. anal.)
 IT 150749-57-8

August 23, 2007

10/803,667

RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses) (fluorescence and intramol. energy transfer in particles for biochem. anal.)

RN 150749-57-8 HCAPLUS
CN Benzo[thiazolium, 3-(3-(trimethylammonio)propyl)-2-[5-(3-(3-(trimethylammonio)propyl)-2(3H)-benzothiazolylidene)-1,3-pentadienyl]-, tribromide (9CI) (CA INDEX NAME)



● 3 Br-

L80 ANSWER 44 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1991:510002 HCAPLUS Full-text

DOCUMENT NUMBER: 115:110002

TITLE: Nucleic acid fractionation by counter migration capillary electrophoresis

INVENTOR(S): Chin, Allan Michael

PATENT ASSIGNEE(S): Applied Biosystems, Inc., USA

SOURCE: PCT Int. Appl., 52 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|---|------|----------|-----------------|----------|
| WO 9102244 | A1 | 19910221 | WO 1990-054380 | 19900806 |
| W: JP | | | | |
| RW: AT, BE, CH, DE, DK, ES, FR, GB, IT, LU, NL, SE | | | | |
| US 5096554 | A | 19920317 | US 1989-390631 | 19890807 |
| US 5110424 | A | 19920505 | US 1990-562790 | 19900806 |
| EP 486559 | A1 | 19920527 | EP 1990-912127 | 19900806 |
| EP 486559 | B1 | 19960320 | | |
| R: AT, BE, CH, DE, DK, ES, FR, GB, IT, LI, LU, NL, SE | | | | |
| JP 04507001 | T | 19921203 | JP 1990-511646 | 19900806 |
| JP 07097101 | B | 19951018 | | |
| AT 135606 | T | 19960415 | AT 1990-912127 | 19900806 |
| PRIORITY APPLN. INFO.: | | | | |
| WO 1990-054380 A 19890807 W 19900806 | | | | |

AB The title method is based on counter migration of nucleic acid fragments in an upstream direction through a polymer-containing (e.g. cellulose derivative-containing) solution which is moving by electroosmotic flow in a downstream direction. Fractionation of selected-size nucleic acid fragments can be enhanced by reducing the difference between the electroosmotic flow rate and the migration rates of the selected-size fragments. An intercalating agent

115

10/803,667

August 23, 2007

may be added to double-stranded fragments to increase preferentially the migration roles of smaller mol. weight fragments through the polymer solution. Schematic diagrams of the electrophoretic system are included, as are electropherograms of fractionated DNA fragments, e.g. a mixture formed by HaeIII digestion of ϕ phi.X174 phage.

IC ICM G01N027-26

CC ICS B01D057-02

IT Virus, Bacterial

IT (phi X174, DNA fragments of, counter migration capillary electrophoresis of fractionation of)

IT 65-61-2, Acridine orange 1239-45-8 107091-89-4, Thiazole orange

RL: ANST (Analytical study, (in counter migration capillary electrophoresis of nucleic acid fragments)

IT 107091-89-4, Thiazole orange

RL: ANST (Analytical study, (in counter migration capillary electrophoresis of nucleic acid fragments)


RN 107091-89-4 HCAPLUS

CN Quinolium, 1-methyl-4-[(3-methyl-2(3H)-benzothiazolylidene)methyl]-, 4-methylbenzenesulfonate (1:1) (CA INDEX NAME)

CM 1

CRN 24144-08-9


CMF C19 H17 N2 S



CM 2

CRN 16722-51-3

CMF C7 H7 O3 S



L80 ANSWER 45 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN

116

August 23, 2007

10/803,667

August 23, 2007

10/803,667

ACCESSION NUMBER: 1990:232277 HCAPLUS Full-text
DOCUMENT NUMBER: 112.232277
TITLE: Fluorescent nucleic acid dye method for multi-parameter flow-cytometric analysis of cellular components of a body fluid
INVENTOR(S): Loken, Michael R.; Terstappen, Leon W. M. M.
PATENT ASSIGNEE(S): Becton, Dickinson and Co., USA
SOURCE: Eur. Pat. Appl., 12 pp.
CODEN: EPXXDW
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|---|------|----------|-----------------|----------|
| EP 347210 | A2 | 19891220 | EP 1989-306036 | 19890614 |
| EP 347210 | A3 | 19901017 | | |
| EP 347210 | B1 | 19940907 | | |
| R: AT, BE, CH, DE, ES, FR, GB, GR, IT, LI, LU, NL, SE | | | | |
| US 5047321 | A | 19910910 | US 1988-207099 | 19880615 |
| AU 8935961 | A | 19891221 | AU 1989-35961 | 19890601 |
| AU 613137 | B2 | 19910725 | | |
| CA 1340170 | C | 19881208 | CA 1989-601444 | 19890601 |
| NO 8902302 | A | 19891218 | NO 1989-2302 | 19890605 |
| NO 175506 | B | 19940711 | | |
| NO 175506 | C | 19941019 | | |
| FI 8902926 | A | 19891216 | FI 1989-2926 | 19890614 |
| FI 94180 | B | 19950413 | | |
| FI 94180 | C | 19950725 | | |
| ES 2063820 | T3 | 19950116 | ES 1989-306036 | 19890614 |
| DK 8902968 | A | 19891216 | DK 1989-2968 | 19890615 |
| JP 02073157 | A | 19900313 | JP 1989-153564 | 19890615 |
| JP 07026954 | B | 19950329 | | |

PRIORITY APPLN. INFO.:
AB The title method uses (1) >1 fluorescence measurements with 22 nucleic acid dyes and 21 fluorescently labeled cell-surface marker, and (2) >1 light scattering measurements. A kit containing the nucleic acid dyes and cell-surface marker is also described. The method of the invention was used in differential blood cell anal. of a thrombocytopenic B-chronic lymphocytic leukemia patient. A blood sample from the patient was incubated with LDS-751 and Thiazole-Orange (nucleic acid dyes) and HLe-1 (anti-CD45 antigen monoclonal antibody) conjugated to phycoerythrin. Frequency of cells occurring in the light-scattering region typical for platelets was low compared with normal peripheral blood, but discrimination of platelets from reticulocytes was still excellent. Separation of leukocytes and reticulocytes was less clear, but these 2 cell types could be clearly separated on the basis of expression of CD45 antigen. Dot plots for the flow-cytometric differential anal. are given. Differential anal. of normal bone marrow cells is also described.


IC ICM G01N033-58
ICS G01N033-50
CC 9-5 (Biochemical Methods)
IT Blood analysis
Cerebrospinal fluid
Peritoneal fluid
Urine analysis
(flow cytometry in, nucleic acid dyes and fluorescence-labeled cell-surface marker for)
IT 76433-29-9, LDS 751 107091-89-4, Thiazole-Orange

117

RL: ASST (Analytical study)
(for differential nucleic acid dye, in flow cytometry of body fluid with fluorescence-labeled cell-surface marker)
IT 76433-29-9, LDS 751 107091-89-4, Thiazole-Orange
RL: AUST (Analytical study)
(for differential nucleic acid dye, in flow cytometry of body fluid with fluorescence-labeled cell-surface marker)
RN 76433-29-9 HCAPLUS
CN Benzothiazolium, 2-[4-[4-(dimethylamino)phenyl]-1,3-butadien-1-yl]-3-ethyl-, perchlorate (1:1) (CA INDEX NAME)

CM 1

CRN 76433-28-8
CMF C21 H23 N2 S



CM 2

CRN 14797-73-0
CMF C1 O4




RN 107091-89-4 HCAPLUS
CN Quinolinium, 1-methyl-4-((3-methyl-2(3H)-benzothiazolylidene)methyl)-, 4-methylbenzenesulfonate (1:1) (CA INDEX NAME)

CM 1


CRN 24144-08-9
CMF C19 H17 N2 S

118



CM 2

CRN 16722-51-3
CMF C7 H7 O3 S



L80 ANSWER 46 OF 46 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 1990:232276 HCAPLUS Full-text
DOCUMENT NUMBER: 112:232276

TITLE: Flow-cytometric method using a nucleic acid dye and optical immunofluorescence for discriminating between intact and damaged cells in a body fluid
Terstappen, Leon W. M. M.; Loken, R. Michael; Shah, Virendra O.

INVENTOR(S): Becton, Dickinson and Co., USA
Eur. Pat. Appl., 14 pp.

PATENT ASSIGNEE(S): Becton, Dickinson and Co., USA

SOURCE: Eur. Pat. Appl., 14 pp.

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|------------|-------------|--|-----------------|----------------|
| EP 347039 | A2 | 19891220 | EP 1989-304754 | 19890510 |
| EP 347039 | A3 | 19901024 | | |
| EP 347039 | B1 | 19931118 | | |
| | R: | AT, BE, CH, DE, ES, FR, GB, GR, IT, LI, LU, NL, SE | | |
| | US 5057413 | A | 19911015 | US 1988-206454 |
| | AT 97500 | T | 19931215 | AT 1989-304754 |
| | ES 2061987 | T3 | 19941216 | ES 1989-304754 |
| | JP 02103464 | A | 19900416 | JP 1989-150348 |
| | | | US 1988-206454 | A 19880613 |
| | | | EP 1989-304754 | A 19890510 |

PRIORITY APPLN. INFO.:

AB The title method uses the nucleic acid dye to selectively stain intact vs. damaged cells, which then may be counted and sorted by flow cytometry. The method may also be used in conjunction with fluorescently labeled monoclonal

antibodies (MAbs) to simultaneously identify cellular antigens. A kit containing the nucleic acid dye and 21 MAbs is described. The method was used to determine the percent of intact nucleated cells in NH₄Cl-lysed, paraformaldehyde-fixed peripheral blood leukocytes from 20 donors, using LDS-751 as the nucleic acid dye. The mean nos. of intact nucleated cells, intact lymphocytes, intact monocytes, intact neutrophils, and intact eosinophils were 34, 88, 56, 76, and 41%, resp. Use of MAbs to a variety of fluorescently-labeled antigens, e.g. phycoerythrin-labeled CD5 and FITC-labeled CD20, in the above method is described.

IC ICM G01N033-58

ICS C12Q001-02; G01N021-75

ICA G01N033-569; G01N033-577

CC 9-5 (Biochemical Methods)

IT Blood analysis

Cerebrospinal fluid

Peritoneal fluid

Urine analysis

dye (flow cytometry in, intact and damaged cell determination by, nucleic acid for)

IT 76433-29-9, LDS 751

RL: ANST (Analytical study)
(as nucleic acid dye, in flow cytometry of intact and damaged cells)

IT 76433-29-9, LDS 751

RL: ANST (Analytical study)

(as nucleic acid dye, in flow cytometry of intact and damaged cells)

RN 76433-29-9 HCAPLUS

CN Benzo[h]azolum, 2-[4-[4-(dimethylamino)phenyl]-1,3-butadien-1-yl]-3-ethyl-, perchlorate (1:1) (CA INDEX NAME)

CM 1

CRN 76433-28-8

CMF C21 H23 N2 S



CM 2

CRN 14797-73-0

CMF Cl O4



L59 50 SEA FILE=HCAPLUS ABB=ON PLU=ON L57 AND L58
 L60 11 SEA FILE=HCAPLUS ABB=ON PLU=ON L56 AND L59
 L61 11 SEA FILE=HCAPLUS ABB=ON PLU=ON L47 AND L52
 L62 11 SEA FILE=HCAPLUS ABB=ON PLU=ON L47 AND L52
 L63 11 SEA FILE=HCAPLUS ABB=ON PLU=ON L47 OR L62
 L64 6 SEA FILE=HCAPLUS ABB=ON PLU=ON L64 AND L57
 L65 5 SEA FILE=HCAPLUS ABB=ON PLU=ON L64 AND L58
 L66 22 SEA FILE=HCAPLUS ABB=ON PLU=ON L65 OR L66 OR L60
 L67 29 SEA FILE=HCAPLUS ABB=ON PLU=ON (L57 OR L58) AND ?BACTER?
 L68 46 SEA FILE=HCAPLUS ABB=ON PLU=ON L67 OR L68
 L69 29 SEA FILE=HCAPLUS ABB=ON PLU=ON L69 AND ?STAIN?
 L70 46 SEA FILE=HCAPLUS ABB=ON PLU=ON L69 OR L71
 L71 4829 SEA FILE=HCAPLUS ABB=ON PLU=ON SAKAI Y7/AU
 L72 2302 SEA FILE=HCAPLUS ABB=ON PLU=ON KAWASHIMA Y7/AU
 L73 989 SEA FILE=HCAPLUS ABB=ON PLU=ON INOUE J7/AU
 L74 293 SEA FILE=HCAPLUS ABB=ON PLU=ON IKEUCHI Y7/AU
 L75 7 SEA FILE=HCAPLUS ABB=ON PLU=ON (L74 OR L75 OR L76 OR L77)
 L76 2 AND ?BACTER? AND ?STAIN?
 L77 2 SEA FILE=HCAPLUS ABB=ON PLU=ON L73 AND L78
 L78 5 SEA FILE=HCAPLUS ABB=ON PLU=ON L78 NOT L79
 L79
 L81

=> d 181 ibib abs tot

L81 ANSWER 1 OF 5 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2005:325608 HCAPLUS Full-text
 DOCUMENT NUMBER: 142:351689

TITLE: Apparatus and method for analyzing bacteria

INVENTOR(S): Kawashima, Yasuyuki
 PATENT ASSIGNEE(S): Sysmex Corporation, Japan
 U.S. Pat. Appl. Publ., 22 pp.

SOURCE: CODEN: USXWCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|---------------|------|----------|-----------------|------------|
| US 2005079569 | A1 | 20050414 | US 2004-961734 | 20041008 |
| JP 2005110629 | A | 20050428 | JP 2003-352170 | 20031010 |
| | | | JP 2003-352170 | A 20031010 |

PRIORITY APPLN. INFO.:
 AB An apparatus for analyzing bacteria is described that includes an analytic sample preparation section for preparing an analytic sample by treating a specimen so as to generate a morphol. difference between Gram-neg. bacteria and Gram-pos. bacteria, a detector for detecting optical information from each particle contained in the analytic sample and an analyzing section for detecting Gram-pos. bacteria contained on the basis of the detected optical information. A method for analyzing bacteria is also described.

L81 ANSWER 2 OF 5 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2004:834918 HCAPLUS Full-text

TITLE: Methods for measuring bacteria,

bacteria measuring apparatuses, and storage media for storing computer-executable programs for analyzing bacteria

INVENTOR(S): Kawashima, Yasuyuki

PATENT ASSIGNEE(S): Sysmex Corporation, Japan

SOURCE: Eur. Pat. Appl.

CODEN: EPXXDW

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|------------|------|----------|-----------------|----------|
| EP 1466985 | A2 | 20041013 | EP 2004-8637 | 20040408 |
| EP 1466985 | A3 | 20041103 | | |
| EP 1466985 | B1 | 20060621 | | |

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK, HR

JP 2004305173 A 20041104 JP 2003-106569 A 20030410

US 2004219627 A 20041104 US 2004-821732 20040408

PRIORITY APPLN. INFO.: JP 2003-106569 A 20030410

AB Methods for measuring bacteria are described that include (a) fluorescently staining bacteria in a sample; (b) detecting size information from the bacteria in the sample, and fluorescence information expressing intensity of fluorescent light emitted by the bacteria; (c) creating a scattergram representing a distribution of the bacteria based on the size information and the fluorescence information detected; (d) analyzing the distribution of the bacteria in the scattergram; and (e) determining whether the bacteria in the sample is bacillus or coccus based on a result of the analyzing. Bacteria measuring apparatuses and storage media for storing computer-executable programs for analyzing bacteria are also described.

L81 ANSWER 3 OF 5 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2004:348049 HCAPLUS Full-text
 DOCUMENT NUMBER: 140:317640

TITLE: Sample analyzers, bacteria analyzers, and solutions for diluting and cleaning

INVENTOR(S): Kawashima, Yasuyuki; Ikeda, Masayuki

PATENT ASSIGNEE(S): Sysmex Corporation, Japan

SOURCE: Eur. Pat. Appl., 40 pp.

CODEN: EPXXDW

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|------------|------|----------|-----------------|----------|
| EP 1413889 | A2 | 20040428 | EP 2003-24425 | 20031023 |
| EP 1413889 | A3 | 20040602 | | |
| EP 1413889 | B1 | 20061220 | | |

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK

JP 2004144633 A 20040520 JP 2002-310585 20021025

JP 3863087 B2 20061227

AT 349014 T 20070115

US 2004096931 A1 20040520

PRIORITY APPLN. INFO.: AT 2003-24425 20031023

US 2003-692554 20031024

JP 2002-310585 A 20021025

AB Sample analyzers for analyzing a sample are described that include a pipet for suctioning the sample; a sample preparation unit for preparing a measured sample by diluting the sample supplied by the pipet with an acidic solution; a pipet washing unit for washing the pipet with the acidic solution; a detection unit for obtaining a detection signal from the measured sample prepared by the sample preparation unit; and a controller for calculating an anal. result from

the detection signal obtained by the detection unit. Bacteria analyzers for analyzing bacteria and solns. for use in sample analyzers are also described.

L81 ANSWER 4 OF 5 HCAPLUS COPYRIGHT 2007 ACS ON STN
 2004:282738 HCAPLUS Full-Text
 Bacteria counting method, bacteria
 counting apparatus, and reagent kit for counting
 bacteria

INVENTOR(S):
 Kawashima, Yasuyuki; Ikeuchi,
 Yoshino; Sakai, Yasuhiro
 PATENT ASSIGNEE(S):
 Sysmex Corporation, Japan
 SOURCE:
 Eur. Pat. Appl.
 CODEN: EPXDXW

DOCUMENT TYPE:
 Patent
 LANGUAGE:
 English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|--|------|----------|-----------------|------------|
| EP 1405918 | A1 | 20040407 | EP 2003-22247 | 20031001 |
| EP 1405918 | B1 | 20070228 | | |
| R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK | | | | |
| JP 2004121143 | A | 20040422 | JP 2002-292606 | 20021004 |
| AT 355387 | T | 20060315 | AT 2003-22247 | 20031001 |
| US 2004067548 | A1 | 20040408 | US 2003-679146 | 20031003 |
| PRIORITY APPLN. INFO.: | | | JP 2002-292606 | A 20021004 |

AB Methods for counting bacteria are described that include: (a) preparing an assay sample by staining a specimen using a fluorescent dye, thereby producing a difference in fluorescent intensity between live bacteria and dead bacteria; (b) detecting optical information from the assay sample; and (c) classifying and counting the live bacteria and the dead bacteria based on the detected optical information. Bacteria counting apparatuses and reagent kits for counting bacteria are also described.

L81 ANSWER 5 OF 5 HCAPLUS COPYRIGHT 2007 ACS ON STN
 2001:100146 HCAPLUS Full-Text
 134:271173

DOCUMENT NUMBER:
 134:271173
 TITLE:
 Formulation design of ointment base suitable for
 healing of lesions in treatment of bedsores

AUTHOR(S):
 Shigezawa, Masato; Ohgaya, Toyooki; Takeuchi,
 Hirofumi; Hino, Tomoaki; Kawashima, Yoshiaki

CORPORATE SOURCE:
 Department of Pharmacy, Takayama Red Cross Hospital,
 Gifu, 506-8550, Japan

SOURCE:
 Chemical & Pharmaceutical Bulletin (2001), 49(2),
 129-133

CODEN: CPBTAL; ISSN: 0009-2363
 PUBLISHER:
 Pharmaceutical Society of Japan
 DOCUMENT TYPE:
 Journal
 LANGUAGE:
 English

AB We intended to develop a desired ointment base suitable for treatment of bedsores including the proliferation of granulation and epidermis. The main bedsores bacteria detected in our hospital were *S. aureus* in gram-pos. coccus and *P. aeruginosa* in gram-neg. bacillus. As the macrogol ointment (MO) was found to have bactericidal effects on these bacteria, MO was adopted as the base for the objective ointment. To improve the properties of the ointment base such as regulating the humidity of the exudation and controlling the

release of antibiotics formulated in the ointment, co-formulating effects of various additives to MO were evaluated. The sustained release function of the ointment base was obtained by adding hydrophilic petrolatum (HP) to MO.

However, the resultant ointment was found to have a poor humidity regulating property. On the other hand, MO containing 5% of hydroxypropyl cellulose (HPC) showed both the humidity regulating and the controlled drug releasing properties. It was considered that HPC particles dispersed in the ointment could be swelled by absorbing water to form a gel network. The curd tension meter tests for the ointments prepared with the various polymers showed that the MO-HPC base, which showed the highest sustained drug releasing property, was found to have the highest hardness. This result means that HPC formulated into the base forms the most rigid gel structure to resist the erosion of the ointment and to control the drug release.

REFERENCE COUNT: 14 THERE ARE 14 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> d his nofil

(FILE 'HOME' ENTERED AT 14:19:46 ON 23 AUG 2007)

FILE 'REGISTRY' ENTERED AT 14:19:55 ON 23 AUG 2007

L1 3608 SEA ABB-ON PLU-ON NC4-C6/ES AND C6/ES AND N-2 AND NR-3 AND C-23

L2 0 SEA ABB-ON PLU-ON L1 AND CLO4/MF

L3 368 SEA ABB-ON PLU-ON L1 AND NC=2

L4 44 SEA ABB-ON PLU-ON L3 AND CL=1 AND O=4

L5 E C23H28N2.CLO4/MF

3 SEA ABB-ON PLU-ON C23H28N2.CLO4/MF

D SCA

E C23H28N2/MF

L6 6 SEA ABB-ON PLU-ON C23H28N2/MF AND L1

D SCA

E BENZENAMINE, 4-(4-(2,3-DIHYDRO-1,3,3-TRIMETHYL-1H-INDOL-2-YL)THYL-1H-INDOL-2-YL)-1,3-BUTADIENYL)-N,N-DIMETHYL-"/CN

D

L8 0 SEA ABB-ON PLU-ON 54268-89-2/CN

FILE 'REGISTRY' ENTERED AT 14:28:55 ON 23 AUG 2007

STR 54268-89-2

L9 1 SEA FAM FUL L9

D SCAN

E C23H27N2.CLO4/MF

E C23H27N2/MF

L11 47 SEA ABB-ON PLU-ON C23H27N2/MF

L12 1 SEA ABB-ON PLU-ON L11 AND L1

D SCA

D

FILE 'REGISTRY' ENTERED AT 14:31:26 ON 23 AUG 2007

STR 76433-26-6

L13 7 SEA FAM FUL L13

D SCA

E C23H27N2.CLO4/MF

E C23H27N2.CLO4

E C23H27N2.CLO4/MF

L15 12 SEA ABB-ON PLU-ON C23H27N2.CLO4/MF

L16 1 SEA ABB-ON PLU-ON L15 AND L1

D SCA

August 23, 2007

10/803,667

L17 FILE 'CAPLUS' ENTERED AT 14:33:31 ON 23 AUG 2007
29 SEA ABB-ON PLU-ON L16

L18 FILE 'STNGUIDE' ENTERED AT 14:33:50 ON 23 AUG 2007

L19 FILE 'REGISTRY' ENTERED AT 14:43:49 ON 23 AUG 2007
E C21H23N2S.C1O4/MF
13 SEA ABB-ON PLU-ON C21H23N2S.C1O4/MF
1 SEA ABB-ON PLU-ON L18 AND NRS=2 AND NR=3 AND NCSC2-C6/ES AND C6/ES
D SCA
E C21H23N2S.C1O4/MF
L*** DEL 13 S E3
E C31H43N4S2.3BR/MF
L20 1 SEA ABB-ON PLU-ON C31H43N4S2.3BR/MF
D SCA
E C53H62N4S2.3BR/MF
E C53H62N6S2.4I/MF
L21 2 SEA ABB-ON PLU-ON C53H62N6S2.4I/MF
L22 2 SEA ABB-ON PLU-ON L21 AND NRS=4 AND NR=8
D SCA
E "QUINOLINUM, 1,1'-(1,3-PROPADEDIYL)BIS"/CN
E "QUINOLINUM, 1,1'-(1,3-PROPADEDIYL)BIS((DIMETHYLIMINO)-3,1-P
E "QUINOLINUM, 1,1'-(1,3-PROPADEDIYL)BIS((DIMETHYLIMINO)-3,1-P
E "QUINOLINUM, 1,1'-(1,3-PROPADEDIYL)BIS((DIMETHYLIMINO)-3,1-P
L23 1 SEA ABB-ON PLU-ON "QUINOLINUM, 1,1'-(1,3-PROPADEDIYL)BIS((DIM
ETHYLIMINO)-3,1-PROPADEDIYL)BIS(4-(3-(3-METHYL-2(3H)-BENZOXAZ
OLYLIDENE)-1-PROPYNYL)-, TETRAIODIDE"/CN
D L21
L24 0 SEA ABB-ON PLU-ON L23 AND L21
D L21
L25 1 SEA ABB-ON PLU-ON L21 NOT 177597-81-8
D SCA
E C26H31N3S.2I/MF
L26 1 SEA ABB-ON PLU-ON C26H31N3S.2I/MF
D SCA

FILE 'STNGUIDE' ENTERED AT 14:54:34 ON 23 AUG 2007

FILE 'REGISTRY' ENTERED AT 15:01:53 ON 23 AUG 2007

L27 E C27H30N4O6S2.2C6H15N/MF
49 SEA ABB-ON PLU-ON C27H30N4O6S2/MF
L28 0 SEA ABB-ON PLU-ON L27 AND NCNC3/ES AND N2C3/ES AND C6/ES
L29 4535 SEA ABB-ON PLU-ON NCNC3/ES AND N2C3/ES AND C6/ES AND NR=3
L30 17 SEA ABB-ON PLU-ON L29 AND S=2 AND O=6 AND N=4
L31 14 SEA ABB-ON PLU-ON L30 AND NC=1
D SCA

L32 E C27H32N4O6S2.C6H15N/MF
2 SEA ABB-ON PLU-ON C27H32N4O6S2.2C6H15N/MF
D SCA

L33 E C35H27BF6N3O7S2.NA/MF
1 SEA ABB-ON PLU-ON C35H27BF6N3O7S2.NA/MF
D SCA

L34 E C35H28BF6N4O7S.NA/MF
1 SEA ABB-ON PLU-ON C35H28BF6N4O7S.NA/MF
D SCA

L35 0 SEA SSS SAM L35
L36 STR
L37 0 SEA SSS FUL L35
L38 STR L35

127

August 23, 2007

10/803,667

L39 2 SEA SSS SAM L38
D SCA
L40 20 SEA SSS FUL L38
L41 10 SEA ABB-ON PLU-ON L40 AND NC=2
L42 STR
L43 1 SEA SSS SAM L42
D SCA
L44 95 SEA SSS FUL L42
L45 48 SEA ABB-ON PLU-ON L44 AND NC=2
L46 67 SEA ABB-ON PLU-ON L16 OR L19 OR L20 OR L25 OR L26 OR L32 OR L33 OR L34 OR L41 OR L45

FILE 'CAPLUS' ENTERED AT 16:33:40 ON 23 AUG 2007
526 SEA ABB-ON PLU-ON L46
E US2004-803667/APPS
L48 1 SEA ABB-ON PLU-ON US2004-803667/AP
SEL RN

FILE 'REGISTRY' ENTERED AT 16:34:32 ON 23 AUG 2007
57 SEA ABB-ON PLU-ON (7803-49-8/BI OR 10182-91-9/BI OR 10182-92-0/BI OR 107-35-7/BI OR 107-95-9/BI OR 107-96-0/BI OR 108-98-5/B
I OR 110-15-6/BI OR 110-17-8/BI OR 110-17-8/BI OR 1119-97-7/BI OR 121-57-3/BI
OR 1310-73-2/BI OR 1333-74-0/BI OR 13881-91-9/BI OR 14797-65-0/
BI OR 15053-09-5/BI OR 150749-57-8/BI OR 15461-40-2/BI OR
157199-63-8/BI OR 166196-17-4/BI OR 189148-50-3/BI OR 24147-36-2/BI OR 335080-22-3/BI OR 33669-61-3/BI OR 361544-71-0/BI OR
361544-72-1/BI OR 50-21-5/BI OR 50-81-7/BI OR 52-90-4/BI OR
5329-14-6/BI OR 56-40-6/BI OR 56-84-8/BI OR 56-85-9/BI OR
56-86-0/BI OR 57-13-6/BI OR 60-24-2/BI OR 60-32-2/BI OR
63-68-3/BI OR 63-74-1/BI OR 6303-21-5/BI OR 68-11-1/BI OR
6899-10-1/BI OR 70-18-8/BI OR 70-47-3/BI OR 74-89-5/BI OR
7440-44-0/BI OR 7558-79-4/BI OR 76433-27-7/BI OR 76433-29-9/BI
OR 7647-01-0/BI OR 77-92-9/BI OR 7704-34-9/BI OR 7778-77-0/BI
OR 7782-44-7/BI OR 7782-99-2/BI OR 877-24-7/BI OR 89-65-6/BI)
L50 10 SEA ABB-ON PLU-ON L49 AND L46

FILE 'CAPLUS' ENTERED AT 16:35:00 ON 23 AUG 2007

D SCA L48
E CYTOMETRY+ALL/CT

FILE 'HCAPLUS' ENTERED AT 16:36:16 ON 23 AUG 2007

L51 8194 SEA ABB-ON PLU-ON CYTOMETRY+PFT,NT/CT

FILE 'CAPLUS' ENTERED AT 16:36:27 ON 23 AUG 2007

D SCA L48

FILE 'HCAPLUS' ENTERED AT 16:36:28 ON 23 AUG 2007

E URINE ANALYSIS+ALL/CT

L52 46411 SEA ABB-ON PLU-ON URINE ANALYSIS+PFT,NT/CT

L53 145 SEA ABB-ON PLU-ON L47 AND (L51 OR L52 OR URIN? OR ?STAIN? OR BLOOD?)

L54 14072 SEA ABB-ON PLU-ON STAINING, BIOLOGICAL+PFT/CT

L55 1842 SEA ABB-ON PLU-ON STAINS, BIOLOGICAL+PFT/CT

L56 58 SEA ABB-ON PLU-ON L47 AND (L54 OR L55)

L*** DEL 0 S L46(L)ANST+NT/CT

L57 192 SEA ABB-ON PLU-ON L46(L)ANST+NT/RL

L58 122 SEA ABB-ON PLU-ON L46(L)BIOL+NT/RL

L59 50 SEA ABB-ON PLU-ON L57 AND L58

L60 11 SEA ABB-ON PLU-ON L56 AND L59

L61 11 SEA ABB-ON PLU-ON L47 AND L52

128

10/803,667

August 23, 2007

L62 11 SEA ABB=ON PLU=ON L47 AND URIN?
L63 22 SEA ABB=ON PLU=ON L61 OR L62 OR L60
L64 11 SEA ABB=ON PLU=ON L61 OR L62
L65 6 SEA ABB=ON PLU=ON L64 AND L57
L66 5 SEA ABB=ON PLU=ON L64 AND L58
L67 22 SEA ABB=ON PLU=ON L65 OR L66 OR L60

FILE 'CAPLUS' ENTERED AT 16:43:08 ON 23 AUG 2007
D SCA L48

FILE 'HCAPLUS' ENTERED AT 16:43:08 ON 23 AUG 2007
L68 29 SEA ABB=ON PLU=ON (L57 OR L58) AND ?BACTER?
L69 46 SEA ABB=ON PLU=ON L67 OR L68
L70 1 SEA ABB=ON PLU=ON L69 AND L48
D SCA
L*** DEL 17 S L59 AND ?STAIN?
L71 29 SEA ABB=ON PLU=ON L69 AND ?STAIN?
L72 1 SEA ABB=ON PLU=ON L71 AND L48
D SCA
L73 46 SEA ABB=ON PLU=ON L69 OR L71
L74 4829 SEA ABB=ON PLU=ON SAKAI Y?/AU
L75 2302 SEA ABB=ON PLU=ON KAWASHIMA Y?/AU
L76 989 SEA ABB=ON PLU=ON INOUE J?/AU
L77 293 SEA ABB=ON PLU=ON KREUCHI Y?/AU
L78 7 SEA ABB=ON PLU=ON (L74 OR L75 OR L76 OR L77) AND ?BACTER?
AND ?STAIN?

FILE 'CAPLUS' ENTERED AT 16:46:42 ON 23 AUG 2007

FILE 'HCAPLUS' ENTERED AT 16:46:51 ON 23 AUG 2007
L79 2 SEA ABB=ON PLU=ON L73 AND L78
L80 46 SEA ABB=ON PLU=ON L73 OR L79
L81 5 SEA ABB=ON PLU=ON L78 NOT L79

FILE 'HCAPLUS' ENTERED AT 16:47:30 ON 23 AUG 2007
D QUE L80
D L80 IBIB ABS HITIND HITSTR TOT
D QUE L81
D L81 IBIB ABS TOT